

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

LEA 33 187

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/744703

INTERNATIONAL APPLICATION NO.

INTERNATIONAL FILING DATE

PRIORITY DATE CLAIMED

PCT/EP99/05073

16 July 1999 (16.07.99)

29 July 1998 (29.07.98)

TITLE OF INVENTION

SUBSTITUTED PYRAZOLE DERIVATIVES

APPLICANT(S) FOR DO/EO/US STRAUB, Alexander; FEURER, Achim; ALONSO-ALJA, Cristina; STAHL, Elke;
STASCH, Johannes-Peter; PERZBORN, Elisabeth; HUTTER, Joachim; DEMBOWSKY, Klaus;

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
- a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
- b. ☐ has been transmitted by the International Bureau.
- c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
- a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
- b. ☐ have been transmitted by the International Bureau.
- c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
- d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A FIRST preliminary amendment.
- ☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:
- 1) Certificate of Mailing under 37 C.F.R. 1.10;
- 2) Transmittal of Information Disclosure Statement under 37 C.F.R. 1.97(b);
- 3) Information Disclosure Citation (Modified Form PTO-1449) and references cited therein; and
- 4) Return Receipt Post Card.

Date of Deposit: 29 January 2001
Express Mail Label No. EK662543167US

- 17.
- ☒
- The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO \$970.00

International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO \$840.00

USPTO international preliminary examination fee (37 CFR 1.482) not paid to USPTO but
international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$690.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$670.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
and all claims satisfied provisions of PCT Article 33(1)-(4) \$96.00

ENTER APPROPRIATE BASIC FEE AMOUNT =**CALCULATIONS PTO USE ONLY**

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total claims	18 - 20 =	0	X \$18.00	\$ 0.00
Independent claims	13 - 3 =	10	X \$78.00	\$ 800.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			0	+ \$260.00
				\$ 0.00

TOTAL OF ABOVE CALCULATIONS = \$ 1,660.00

Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement
must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

SUBTOTAL = \$ 1,660.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)).

TOTAL NATIONAL FEE = \$ 1,660.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +

TOTAL FEES ENCLOSED = \$ 1,660.00

Amount to be
refunded: \$
charged: \$

- a. ☐ A check in the amount of \$ _____ to cover the above fees is enclosed.
- b. ☒ Please charge my Deposit Account No. 13-3372 in the amount of \$ 1,660.00 to cover the above fees.
A duplicate copy of this sheet is enclosed.
- c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
overpayment to Deposit Account No. 13-3372. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO

Jeffrey M. Greenman
Vice President, Patents and Licensing
Bayer Corporation
400 Morgan Lane
West Haven, CT 06516

Jerrie L. Chin
SIGNATURE

Jerrie L. Chin

NAME

41,670

REGISTRATION NUMBER

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Straub, et al.

Serial No.: National Stage Filing of PCT/EP99/05073

Filed: herewith

For: Substituted Pyrazole Derivatives

BOX PCT
Assistant Commissioner for Patents
Washington, D.C. 20231

CERTIFICATE OF MAILING UNDER 37 CFR 1.10

I hereby certify that the *attached* correspondence comprising:


- Transmittal Letter to the United States Designated/Elected Office (DO/EO/US) Concerning a Filing under 35 U.S.C. 371 [IN DUPLICATE];
- Original Oath or Declaration of the Inventor(s) (35 U.S.C. 371(c)(4));
- English Translation of the Application (35 U.S.C. 371(c)(2));
- Copy of the International Application as filed (35 U.S.C. 371(c)(2));
- Information Disclosure Statement under 37 C.F.R. 1.97 and 1.98 consisting of Transmittal of Information Disclosure Statement under 37 C.F.R. 1.97(b), Information Disclosure Citation (Modified Form PTO-1449) and copies of references cited therein; and
- Return Receipt Post Card.

is, on the date shown below, being deposited with the United States Postal Service, in an envelope as "Express Mail Post Office to Addressee" Mailing Label Number EK662543167US, addressed to:

Box PCT
Assistant Commissioner for Patents
Washington, D.C. 20231

January 29, 2001

Date



Signature of Person Certifying / Beatriz Alviz

PATENT
Atty. Docket No.: Le A 33 187

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICANTS: Straub, *et al.*

SERIAL NO.: 09/744,703

FILING DATE: January 29, 2001

TITLE: Substituted Pyrazole Derivatives

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

This Preliminary Amendment is submitted in the above-captioned application having U.S.S.N. 09/744,703. Please amend the application as follows:

In the Claims

Please cancel claims 9 and 12.

Please amend claims 4, 6, 7, 8 and 13-18 as shown in the attached pages. A marked version of the claim set showing the changes made is also attached.

Please add new claims 19-21 as shown in the attached page.

Remarks

By way of this Preliminary Amendment, claims 1-8, 10, 11 and 13-21 are pending. Claims 9 and 12 have been canceled. Claims 4, 6, 7, 8 and 13-18 have been amended to put the claims in a format appropriate for U.S. prosecution. Claims 19-21 have been added to include material deleted from amended claims 6 and 18. No new matter has been added by way of these amendments.

More specifically, claim 4 has been amended to specify that it is dependent on claim 1. Claim 6 has been amended to specify that R¹, R², R³, R³⁶, R³⁷, R³⁸, R³⁹, R^{36'}, R^{37'}, R^{38'}, R^{39'}, X, Y and A are as defined in Claim 1. Additionally, claim 6 has been amended to remove phrases such as "if appropriate," "preferably," and "customary". Material deleted from Claim 6 has been rewritten in new claims 19 and 20. Claim 8 has been amended to add the recitation of a pharmaceutically acceptable carrier to the claim. Claims 13-18 have been amended to convert the Swiss-type use claim to the U.S. method of treatment format. Finally, claim 18 has been amended so that the claim contains the proper multiple dependent claim format in compliance with 37 C.F.R. § 1.75(c) and to separate the claim into two independent claims. Material deleted from claim 18 has been rewritten in independent form in new claim 21.

Conclusion

Applicants respectfully submit that the pending claims, as amended, are in condition for allowance. Please charge any fees due with this amendment to deposit account number 13-3372. If the Examiner believes that a conversation with Applicants' attorney would be helpful in expediting prosecution of this application, the Examiner is invited to call the undersigned attorney at (203) 812-3964.

Respectfully submitted,

Dated: *March 22, 2001*


Bayer Corporation
400 Morgan Lane
West Haven, CT 06516
(Tel) (203) 812-3964
(Fax) (203) 812-5492
e-mail: jerrie.chiu.b@bayer.com

Jerrie L. Chiu
Jerrie L. Chiu
Attorney for Applicants
Reg. No. 41,670

4. (Amended) Compounds according to claim 1,

in which

R^1 , X and Y are attached to the pyrimidine ring as follows



and

R^1 represents an optionally substituted cyclopropyl, cyclobutyl, cyclopentenyl, cyclopentyl, cyclohexyl, 1-hydroxycyclopropyl or 1-(fluoromethyl)cyclopropyl radical,

X represents NH_2


and

Y represents hydrogen or NH_2 .

6. (Amended) Process for preparing the compounds of the general formula (I) according to Claim 1, characterized in that,

depending on the various meanings of the heterocycles listed above under R^2 and R^3 ,

[A] compounds of the general formula (II)




in which

R¹, X and Y are each as defined above in claim 1,

and

D represents radicals of the formulae



in which

R³⁸ represents C₁-C₄-alkyl


are converted, by reaction with compounds of the general formula (III)



in which

A is as defined above in claim 1,

in inert solvents into the compounds of the general formula (IV) or (IVa)



in which


A, X, Y and R¹ are each as defined above in claim 1,

and, in the case of the compounds of the general formula (IVa), are subsequently cyclized with carboxylic acids, nitriles, formamides or guanidium salts,

and, in the case of the compounds of the general formula (IV), are cyclized with 1,3-dicarbonyl derivatives, their salts, tautomers, enol ethers or enamines, in the presence of acids,

or


[B] in the case that R² and R³ together form a pyrazine ring, compounds of the general formula (IV) are initially converted by nitrosation into the compounds of the general formula (V)



in which

A, X, Y and R¹ are each as defined above in claim 1,

in a second step, the compounds of the general formula (VI)



in which


A, X, Y and R¹ are each as defined above in claim 1,

are prepared by a reduction,

and these are subsequently cyclized with 1,2-dicarbonyl compounds,

or

[C] compounds of the general formula (VII)



in which

A^1 , R^2 and R^3 are each as defined above in claim 1,

and

L represents a radical of the formula $-\text{SnR}^{36}\text{R}^{37}\text{R}^{38}$, ZnR^{39} , iodine, bromine or triflate,


in which

R^{36} , R^{37} and R^{38} are identical or different and each represents straight-chain or branched alkyl having up to 4 carbon atoms

and

R^{39} represents halogen

are reacted with compounds of the general formula (VIII)



in which

X, Y and R¹ are each as defined above in claim 1,

and

in the case that L = SnR³⁶R³⁷R³⁸ or ZnR³⁹,

T represents triflate or represents halogen,

and,

in the case that L = iodine, bromine or triflate,

T represents a radical of the formula SnR³⁶R³⁷R³⁸, ZnR³⁹ or BR⁴⁰R⁴¹,

in which


R³⁶, R³⁷, R³⁸ and R³⁹ have the meanings of R³⁶, R³⁷, R³⁸ and R³⁹ given above in claim 1 and are identical to or different from them,

R⁴⁰ and R⁴¹ are identical or different and each represent hydroxyl, aryloxy having 6 to 10 carbon atoms or straight-chain or branched alkyl or alkoxy having in each case up to 5 carbon atoms, or together form a 5- or 6-membered carbocyclic ring

in a palladium-catalysed reaction in inert solvents,

or

[D] amidines of the general formula (IX)




(IX),

in which

A, R² and R³ are each as defined above in claim 1,


are reacted with compounds of the general formula (X), (Xa), (Xb) or (Xc)




(X)



(Xa)



(Xb)



(Xc) ,

in which

R¹ represents the optionally substituted cycloalkyl radical listed above under R¹;

Alk represents straight-chain or branched alkyl having up to 8 carbon atoms,

and

Z represents an NH₂ group, a monoalkylamino group having up to 7 carbon atoms, a dialkylamino group having up to 7 carbon atoms, a piperidinyl or morpholinyl radical which is attached via the nitrogen, hydroxyl, alkoxy having up to 7 carbon atoms, acyloxy having up to 7 carbon atoms or aryloxy having 6 to 10 carbon atoms,

and the substituents listed under X, Y, R¹, R², R³ and/or A are modified or introduced by acylation and derivatization of free amino groups, chlorination, catalytic hydrogenation, reduction, oxidation, removal of protective groups and/or nucleophilic substitution.

8. (Amended) Medicaments, comprising at least one compound of the general formula (I) according to Claim 1 and a pharmaceutically acceptable carrier.
13. (Amended) A method of treating a cardiovascular disease, comprising administering to a mammal an effective amount of a compound according to Claim 1.
14. (Amended) The method of claim 13, wherein said cardiovascular disease is hypertension.
15. (Amended) A method of treating thromboembolic disorders and ischaemia, comprising administering to a mammal an effective amount of a compound according to claim 1.
16. (Amended) A method of treating sexual dysfunction, comprising administering to a mammal an effective amount of a compound according to claim 1.
17. (Amended) A method of treating inflammation, comprising administering to a mammal an effective amount of a compound according to claim 1.

19. (New) The process of claim 6, wherein said compound of the general formula (IV) further are cyclized with 1,3-dicarbonyl derivatives, their salts, tautomers, enol ethers or enamines in the presence of acids under microwave irradiation.
20. (New) The process of claim 6, wherein said 1,2-dicarbonyl compound is aqueous glyoxal solution.
21. (New) The method of claim 13, 14, 15, 16 or 17, wherein said compound of the general formula (I) according to Claim 1 is administered in combination with a compound that inhibits the degradation of cyclic guanosine monophosphate (cGMP).

09/744,703
Le A 33 187

Version with Markings to Show Changes Made


- 110 -

Claim 1

4. Compounds according to ~~any of the preceding claims,~~

in which

- 5 R^1 , X and Y are attached to the pyrimidine ring as follows



and

10

R^1 represents an optionally substituted cyclopropyl, cyclobutyl, cyclopentenyl, cyclopentyl, cyclohexyl, 1-hydroxycyclopropyl or 1-(fluoromethyl)cyclopropyl radical,

15

X represents NH_2

and

20

Y represents hydrogen or NH_2 .


5. Compounds according to claim 4, in which R^1 represents an optionally substituted cyclopropyl radical.

6. Process for preparing the compounds of the general formula (I) according to Claim 1,
25 characterized in that,

depending on the various meanings of the heterocycles listed above under R^2 and R^3 ,

[A] compounds of the general formula (II)

5



in which


10

R^1 , X and Y are each as defined above *in Claim 1*,

and

D represents radicals of the formulae

15




in which

20

R^{38} represents C_1 - C_4 -alkyl


are converted, by reaction with compounds of the general formula (III)




in which

A is as defined above *in Claim 1*,

in inert solvents ~~if appropriate in the presence of a base~~ into the compounds of the general formula (IV) or (IVa)



(IV) and



(IVa),

in which

A, X, Y and R¹ are each as defined above,


in Claim 1

and, in the case of the compounds of the general formula (IVa), are subsequently cyclized with carboxylic acids, nitriles, formamides or guanidium salts,

and, in the case of the compounds of the general formula (IV), are cyclized with 1,3-dicarbonyl derivatives, their salts, tautomers, ~~enol ethers or enamines~~, in the presence of acids ~~and, if appropriate, under microwave irradiation~~.

or

[B] in the case that R^2 and R^3 together form a pyrazine ring, compounds of the general formula (IV) are initially converted by nitrosation into the compounds of the general formula (V)




in which

A, X, Y and R^1 are each as defined above.

in Claim 1

in a second step, the compounds of the general formula (VI)



in which


A, X, Y and R^1 are each as defined above *in Claim 1,*

are prepared by a reduction,

and these are subsequently cyclized with 1,2-dicarbonyl compounds, ~~preferably~~
~~aqueous glyoxal solution.~~

or

[C] compounds of the general formula (VII)



5

in which

A¹, R² and R³ are each as defined above *in Claim 1*,

10

and

L represents a radical of the formula -SnR³⁶R³⁷R³⁸, ZnR³⁹, iodine, bromine or triflate,

15

in which

R³⁶, R³⁷ and R³⁸ are identical or different and each represents straight-chain or branched alkyl having up to 4 carbon atoms

20


and

R³⁹ represents halogen

are reacted with compounds of the general formula (VIII)

25

09744703 032601
109220 504446



(VIII),

in which

X, Y and R¹ are each as defined above *in Claim 1,*

and

in the case that L = SnR³⁶R³⁷R³⁸ or ZnR³⁹,

T represents triflate or represents halogen, ~~preferably bromine~~

and,

in the case that L = iodine, bromine or triflate,

T represents a radical of the formula SnR³⁶R³⁷R³⁸, ZnR³⁹ or BR⁴⁰R⁴¹,

in which


R³⁶, R³⁷, R³⁸ and R³⁹ have the meanings of R³⁶, R³⁷, R³⁸ and R³⁹ given above and
are identical to or different from them, *in Claim 1*

R⁴⁰ and R⁴¹ are identical or different and each represent hydroxyl, aryloxy
having 6 to 10 carbon atoms or straight-chain or branched alkyl or alkoxy
having in each case up to 5 carbon atoms, or together form a 5- or
6-membered carbocyclic ring

in a palladium-catalysed reaction in inert solvents, ~~if appropriate in the presence~~
~~of a base,~~

or

5 [D] amidines of the general formula (IX)



in which

10

A, R² and R³ are each as defined above *in Claim 1,*

are reacted ~~for example~~ with compounds of the general formula (X), (Xa),
 (Xb) or (Xc)


15




(X)



(Xa)



(Xb)



(Xc)

in which

R^r represents the optionally substituted cycloalkyl radical listed above under R¹;


Alk represents straight-chain or branched alkyl having up to 8 carbon atoms,
~~preferably up to four carbon atoms; A~~

and

Z represents an NH₂ group, a monoalkylamino group having up to 7 carbon atoms, a dialkylamino group having up to 7 carbon atoms, a piperidyl or morpholyl radical which is attached via the nitrogen, hydroxyl, alkoxy having up to 7 carbon atoms, acyloxy having up to 7 carbon atoms or aryloxy having 6 to 10 carbon atoms,

and, if appropriate, the substituents listed under X, Y, R¹, R², R³ and/or A are modified or introduced by ~~customary methods, preferably by~~ acylation and derivatization of free amino groups, chlorination, catalytic hydrogenation, reduction, oxidation, removal of protective groups and/or nucleophilic substitution.

7. Amidines of the general formula (IX)



(IX),

in which

R¹, R² and A are each as defined in one of the preceding Claims 1 to 3.

and their isomeric forms and salts.

- 5 8. Medicaments, comprising at least one compound of the general formula (I) according to Claim 1, *is a pharmaceutically acceptable carrier*
- 10 9. ~~Process for preparing medicaments, characterized in that at least one compound of the formula (I) according to Claim 1, if appropriate with customary auxiliaries and additives, is converted into a suitable administration form.~~ *canceled*
- 15 10. Medicaments, comprising at least one compound of the general formula (I) according to Claim 1 in combination with organic nitrates or NO donors.
- 20 11. Medicaments, comprising at least one compound of the general formula (I) according to Claim 1 in combination with compounds which inhibit the degradation of cyclic guanosine monophosphate (cGMP).
- 25 12. ~~Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments.~~ *canceled*
- 30 13. *A method of treating a*
~~Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments for the treatment of cardiovascular diseases.~~ *comprising administering to a mammal an effective amount of a compound according to Claim 1.*
14. ~~Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments for the treatment of hypertension.~~ *The method of Claim 13, wherein said cardiovascular disease is*
15. ~~Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments for the treatment of thromboembolic disorders and ischaemia.~~ *A method of treating comprising administering to a mammal an effective amount of a compound according to Claim 1*
16. ~~Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments for the treatment of sexual dysfunction.~~ *A method of treating*

09744703-032601

A method of treating inflammation, comprising

17. ~~Use of compounds of the general formula (I) according to Claim 1 for~~
~~administering to a mammal an effective amount~~
~~of a compound according to Claim 1.~~
~~preparing medicament having anti-inflammatory properties.~~

- 5 18. ~~The method of~~ 13, 14, 15, 16 or 17
~~Use according to any of Claims 12 to 17 where the compounds of the general~~
~~formula (I) according to Claim 1 are used in combination with organic nitrates~~
~~or NO donors~~ in said
is administered an
~~of cyclic guanosine monophosphate (cGMP).~~

09744703.032601

JC07 Rec'd PCT/PTO 29 JAN 2001

Substituted pyrazole derivatives

The present invention relates to substituted pyrazole derivatives, to processes for their preparation and to their use as medicaments, in particular as medicaments for the treatment of cardiovascular disorders.


It is already known that 1-benzyl-3-(substituted heteroaryl)-fused pyrazole derivatives inhibit thromocyte aggregation (cf. EP 667 345 A1).

WO 98/16223 discloses the use of 1-benzyl-3-(substituted hetaryl)-fused pyrazole derivatives for the treatment of specific disorders of the cardiovascular system and the central nervous system.

WO 98/16507 discloses heterocyclymethyl-substituted pyrazole derivatives and their use in the treatment of cardiovascular disorders.

WO 98/23619 likewise discloses substituted pyrazole derivatives for the treatment of cardiovascular disorders.

The present invention relates to substituted pyrazole derivatives of the general formula (I)



in which

at least one of the substituents R^1 , X and Y represents saturated or partially unsaturated C_3 - C_8 -cycloalkyl,

which may optionally be mono- or polysubstituted by amino, azido, formyl, mercaptyl, carboxyl, hydroxyl, morpholino, piperidino, pyrrolidino, sulphonamino, straight-chain, cyclic or branched acyl, acylamino, alkoxy, benzyloxy, alkylamino, dialkylamino, alkylsulphonyl, alkylsulphonamino, alkylthio, alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, halogen, phenyl

and/or is optionally substituted by

straight-chain or branched or cyclic alkyl having up to 6 carbon atoms which for its part may be substituted by amino, mercaptyl, carboxyl, hydroxyl, morpholino, piperidino, pyrrolidino, straight-chain, cyclic or branched acyl, acylamino, alkoxy, alkylamino, dialkylamino, alkylsulphonyl, alkylthio, phenyl, alkylsulphonamino, alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, halogen,


and where the optionally remaining radicals R^1 , X and/or Y are identical or different and each represents hydrogen, azido, formyl, mercaptyl, carboxyl, hydroxyl, straight-chain or branched acyl, alkoxy, alkylthio or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, halogen, phenyl or straight-chain or branched alkenyl or alkynyl having in each case up to 6 carbon atoms or alkyl having up to 20 carbon atoms, where both alkenyl, alkynyl and/or alkyl may optionally be substituted by hydroxyl, amino, azido, carboxyl, straight-chain or branched acyl, alkoxy, alkoxycarbonyl or acylamino having in each case up to 5 carbon atoms, aryl having 6 to 10 carbon atoms, a 5- to 6-membered aromatic heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, halogen, cyano, dialkylamino having up to 6 carbon

atoms, alkylamino having up to 6 carbon atoms and/or cycloalkyl having 3 to 8 carbon atoms or by a radical of the formula $-OR^4$,

in which

R^4 represents straight-chain or branched acyl having up to 5 carbon atoms,

and/or the optionally remaining radicals R^1 , X and/or Y each represent a radical of the formula



in which

a, b and b' are identical or different and represent a number 0, 1, 2 or 3,

R^5 represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

and/or the optionally remaining radicals R^1 , X and/or Y each represent a 3- to 8-membered ring, which may also be linked to the pyrimidine via a $-\text{CO}-$ bridge, and which may be saturated, unsaturated and/or partially unsaturated, which may contain 1 to 4 heteroatoms from the group consisting of N, O, S, SO, SO_2 , which may also be attached via N, which may optionally contain a carbonyl group as ring member and which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl, amino, halogen, carboxyl,

straight-chain or branched acyl, alkoxy, alkoxycarbonyl or acylamino having in each case up to 5 carbon atoms,

and/or the optionally remaining radicals R^1 , X and/or Y each represent straight-chain or branched acyl having up to 6 carbon atoms which is optionally substituted by halogen,

or

represent straight-chain or branched acyloxy having up to 6 carbon atoms,

or

represent arylthio having 6 to 10 carbon atoms or heteroarylthio,

and/or the optionally remaining radicals R^1 , X and/or Y represent radicals of the formulae $-SO_3H$ or $S(O)_cR^6$,

in which

c represents a number 1 or 2,

R^6 represents straight-chain or branched alkyl having 1 to 10 carbon atoms, cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or a 5- to 6-membered heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, where the ring systems may optionally be substituted by halogen or by straight-chain or branched alkyl or alkoxy having in each case up to 4 carbon atoms,

and/or the optionally remaining radicals R^1 , X and/or Y each represent a radical of the formula $PO(OR^7)(OR^8)$,

in which

R^7 and R^8 are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 8 carbon atoms or cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or benzyl,

and/or the optionally remaining radicals R^1 , X and/or Y each represents oxycycloalkyl having 3 to 8 carbon atoms or represent radicals of the formulae $-NH-C(=NH)NH_2$, $CON=C(NH_2)_2$ or $-C=NH(NH_2)$, $(CO)_dNR^9R^{10}$ or $-NHCONR^9R^{10}$,

in which


d represents a number 0 or 1,

R^9 and R^{10} are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 14 carbon atoms or cycloalkyl having 3 to 14 carbon atoms, aryl having 6 to 10 carbon atoms or a 3- to 10-membered ring having up to 5 heteroatoms from the group consisting of N, O, S, which may also be attached via N, where the abovementioned radicals may optionally be substituted by aryl having 6 to 10 carbon atoms, heterocyclyl, cycloalkyl having 3 to 7 carbon atoms, hydroxyl, amino or straight-chain or branched alkoxy, acyl or alkoxycarbonyl having in each case up to 6 carbon atoms,

and in the case that $d = 0$,

R^9 and R^{10} may also represent straight-chain, branched or cyclic acyl having up to 14 carbon atoms, straight-chain or branched hydroxyalkyl having up to 6 carbon atoms, straight-chain or branched alkoxyalkyl having a total of up to 12 carbon atoms, straight-chain or branched

alkoxycarbonyl or acyloxyalkyl having in each case up to 6 carbon atoms or a radical of the formula $-\text{SO}_2\text{R}^{11}$ or radicals of the formulae



5


in which


R^{11} represents straight-chain or branched alkyl having up to 4 carbon atoms,

10

and/or

R^9 and R^{10} represent radicals of the formulae





in which

R^{12} , R^{13} and R^{15} to R^{28} are identical or different and each represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

e represents a number 0, 1 or 2,

and

R^{14} represents straight-chain or branched alkyl having up to 6 carbon atoms or cycloalkyl having 3 to 8 carbon atoms,

R^9 and R^{10} are identical or different and each represents hydrogen; alkyl having up to 14 carbon atoms which is optionally substituted by hydroxyl or alkoxy having up to 8 carbon atoms; aryl having 6 to 10 carbon atoms which is optionally substituted by halogen,

R^2 and R^3 form, together with the double bond, a fused phenyl ring or a fused 6-membered saturated or aromatic heterocycle having up to 3 heteroatoms from the group consisting of N, S and O,

which is optionally substituted up to 3 times by identical or different substituents from the group consisting of formyl, carboxyl, hydroxyl, mercaptyl, straight-chain or branched acyl, alkylthio or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, halogen or straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms which for its part may be substituted by hydroxyl, amino, carboxyl, straight-chain or branched acyl, alkoxy or alkoxycarbonyl having in each case up to 5 carbon atoms,

and/or the fused phenyl ring or the fused 6-membered saturated or aromatic heterocycle is optionally substituted by a group of the formula $-NR^{29}R^{30}$,

in which

R^{29} and R^{30} are identical or different and each represents hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,

or

R^{29} represents hydrogen and

5 R^{30} represents acyl having up to four carbon atoms

and/or the fused phenyl ring or the fused 6-membered saturated or aromatic heterocycle are optionally substituted by phenyl which for its part may be substituted up to 2 times by identical or different substituents from the group consisting of halogen, straight-chain or branched alkyl and alkoxy having in each case up to 6 carbon atoms

and/or the fused phenyl ring or the fused 6-membered saturated or aromatic heterocycle are optionally substituted by a group of the formula
15 $-N=CH-NR^{31}R^{32}$,

in which

R^{31} and R^{32} are identical or different and each represents hydrogen, phenyl or straight-chain or branched alkyl having up to 6 carbon atoms,

A represents a 5- or 6-membered aromatic or saturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O or represents phenyl,

25 which are optionally substituted up to 3 times by identical or different substituents from the group consisting of amino, mercaptyl, hydroxyl, formyl, carboxyl, straight-chain or branched acyl, alkylthio, alkoxyacyl, alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, trifluoromethyl, azido, halogen, phenyl and straight-chain or branched alkyl
30 having up to 6 carbon atoms which for its part may be substituted by

hydroxyl, carboxyl, straight-chain or branched acyl, alkoxy or alkoxycarbonyl
having in each case up to 5 carbon atoms,

and/or

is substituted by a group of the formula $-(CO)_f-NR^{33}R^{34}$,

in which

f represents a number 0 or 1,

R^{33} and R^{34} are identical or different and each represents hydrogen,
phenyl, benzyl or straight-chain or branched alkyl or acyl
having in each case up to 5 carbon atoms,

and their isomeric forms and salts.

The compounds of the general formula (I) according to the invention may also be
present in the form of their salts. In general, salts with organic or inorganic bases or
acids may be mentioned here.

In the context of the present invention, preference is given to physiologically
acceptable salts. Physiologically acceptable salts of the compounds according to the
invention may be salts of the substances according to the invention with mineral
acids, carboxylic acids or sulphonic acids. Particular preference is given, for
example, to salts with hydrochloric acid, hydrobromic acid, sulphuric acid,
phosphoric acid, methanesulphonic acid, ethanesulphonic acid, p-toluenesulphonic
acid, benzenesulphonic acid, naphthalenedisulphonic acid, acetic acid, propionic
acid, lactic acid, tartaric acid, citric acid, fumaric acid, maleic acid or benzoic acid.

Physiologically acceptable salts may also be the metal or ammonium salts of the compounds according to the invention which have a free carboxyl group. Particular preference is given, for example, to sodium, potassium, magnesium or calcium salts, and to ammonium salts which are derived from ammonia, or organic amines, such as, for example, ethylamine, di- or triethylamine, di- or triethanolamine, dicyclohexylamine, dimethylaminoethanol, arginine, lysine or ethylenediamine.

The compounds according to the invention may exist in stereoisomeric forms which are either like image and mirror image (enantiomers) or which are not like image and mirror image (diastereomers). The invention relates both to the enantiomers or diastereomers and to their respective mixtures. The racemates, like the diastereomers, can be separated into stereoisomerically uniform components in a known manner.

In the context of the present invention, the substituents have, unless indicated otherwise, generally the following meanings:

Alkyl generally represents a straight-chain or branched hydrocarbon radical having 1 to 20 carbon atoms. Examples which may be mentioned are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, isopentyl, hexyl, isohexyl, heptyl, isoheptyl, octyl and isooctyl, nonyl, decyl, dodecyl, eicosyl.

Alkenyl generally represents a straight-chain or branched hydrocarbon radical having 2 to 20 carbon atoms and one or more, preferably one or two, double bonds. Examples which may be mentioned are allyl, propenyl, isopropenyl, butenyl, isobutenyl, pentenyl, isopentenyl, hexenyl, isohexenyl, heptenyl, isoheptenyl, octenyl, isooctenyl.

Alkynyl generally represents a straight-chain or branched hydrocarbon radical having 2 to 20 carbon atoms and one or more, preferably one or two, triple bonds. Examples which may be mentioned are ethynyl, 2-butynyl, 2-pentynyl and 2-hexynyl.

Acyl generally represents straight-chain or branched lower alkyl having 1 to 9 carbon atoms which is attached via a carbonyl group. Examples which may be mentioned are: acetyl, ethylcarbonyl, propylcarbonyl, isopropylcarbonyl, butylcarbonyl and isobutylcarbonyl.

5


Alkoxy generally represents a straight-chain or branched hydrocarbon radical having 1 to 14 carbon atoms which is attached via an oxygen atom. Examples which may be mentioned are methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, pentoxy, isopentoxy, hexoxy, isohexoxy, heptoxy, isoheptoxy, octoxy or isooctoxy. The terms "alkoxy" and "alkyloxy" are used synonymously.

10

Alkoxyalkyl generally represents an alkyl radical having up to 8 carbon atoms which is substituted by an alkoxy radical having up to 8 carbon atoms.

15

Alkoxycarbonyl can be depicted, for example, by the formula



20

Alkyl here generally represents a straight-chain or branched hydrocarbon radical having 1 to 13 carbon atoms. The following alkoxycarbonyl radicals may be mentioned as examples: methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl or isobutoxycarbonyl.

25

Cycloalkyl generally represents a cyclic hydrocarbon radical having 3 to 8 carbon atoms. Preference is given to cyclopropyl, cyclopentyl and cyclohexyl. Examples which may be mentioned are cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

Cycloalkoxy represents, in the context of the invention, an alkoxy radical whose hydrocarbon radical is a cycloalkyl radical. The cycloalkyl radical generally has up to

8 carbon atoms. Examples which may be mentioned are: cyclopropyloxy and cyclohexyloxy. The terms "cycloalkoxy" and "cycloalkyloxy" are used synonymously.

- 5 Aryl generally represents an aromatic radical having 6 to 10 carbon atoms. Preferred aryl radicals are phenyl and naphthyl.

Halogen represents, in the context of the invention, fluorine, chlorine, bromine and iodine.

10

Heterocycle represents, in the context of the invention, a saturated, unsaturated or aromatic 3- to 10-membered, for example 5- or 6-membered, heterocycle which may contain up to 3 heteroatoms from the group consisting of S, N and O and which, in the case of a nitrogen atom, may also be attached via this nitrogen atom. Examples which
15 may be mentioned are: oxadiazolyl, thiadiazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, thienyl, furyl, pyrrolyl, pyrrolidinyl, piperazinyl, tetrahydropyranlyl, tetrahydrofuranyl, 1,2,3-triazolyl, thiazolyl, oxazolyl, imidazolyl, morpholinyl or piperidyl. Preference is given to thiazolyl, furyl, oxazolyl, pyrazolyl, triazolyl, pyridyl, pyrimidinyl, pyridazinyl and tetrahydropyranlyl. The term
20 "heteroaryl" (or "hetaryl") represents an aromatic heterocyclic radical.

20

Preference is given to compounds of the general formula (I) according to the invention,

25 in which

30

at least one of the substituents R^1 , X and Y represents cyclopropyl, cyclobutyl, cyclopentenyl, cyclopentyl, cyclohexyl or cycloheptyl which may optionally be mono- or polysubstituted by amino, azido, formyl, mercaptyl, carboxyl, hydroxyl, morpholino, piperidino, pyrrolidino, sulphonamino, straight-chain, cyclic or branched acyl, acylamino, alkoxy, benzyloxy, alkylamino,

dialkylamino, alkylsulphonyl, alkylsulphonamino, alkylthio, alkoxycarbonyl having in each case up to 4 carbon atoms, nitro, cyano, halogen, phenyl

and/or is optionally substituted by


straight-chain or branched or cyclic alkyl having up to 4 carbon atoms which for its part may be substituted by amino, mercaptyl, carboxyl, hydroxyl, morpholino, piperidino, pyrrolidino, straight-chain, cyclic or branched acyl, acylamino, alkoxy, alkylamino, dialkylamino, alkylsulphonyl, alkylthio, phenyl, alkylsulphonamino, alkoxycarbonyl having in each case up to 4 carbon atoms, nitro, cyano, halogen,

and where the optionally remaining radicals R^1 , X and/or Y are identical or different and each represents hydrogen, azido, formyl, mercaptyl, carboxyl, hydroxyl, straight-chain or branched acyl, alkoxy, alkylthio or alkoxycarbonyl having in each case up to 4 carbon atoms, nitro, cyano, halogen, phenyl or straight-chain or branched alkenyl or alkynyl having in each case up to 4 carbon atoms or alkyl having up to 18 carbon atoms, where both alkenyl, alkynyl and/or alkyl may optionally be substituted by hydroxyl, amino, azido, carboxyl, straight-chain or branched acyl, alkoxy, alkoxycarbonyl or acylamino having in each case up to 4 carbon atoms, phenyl, naphthyl or pyridyl, halogen, cyano, dialkylamino having up to 6 carbon atoms, alkylamino having up to 4 carbon atoms and/or cyclopropyl, cyclopentyl, cyclohexyl or by a radical of the formula $-OR^4$,

in which

R^4 represents straight-chain or branched acyl having up to 4 carbon atoms,

and/or the optionally remaining radicals R¹, X and/or Y each represent a radical of the formula



5

in which

a, b and b' are identical or different and represent a number 0, 1 or 2,

10

and/or the optionally remaining radicals R¹, X and/or Y each represent a 3- to 8-membered ring, which may also be linked to the pyrimidine via a -CO-bridge, and which may be saturated, unsaturated and/or partially unsaturated, which may contain 1 to 3 heteroatoms from the group consisting of N, O, S, SO, SO₂, which may also be attached via N, which may optionally contain a carbonyl group as ring member and which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched alkyl having up to 4 carbon atoms which is optionally substituted by hydroxyl, amino, halogen, carboxyl, straight-chain or branched acyl, alkoxy, alkoxy-carbonyl or acylamino having in each case up to 4 carbon atoms,

15

20

and/or the optionally remaining radicals R¹, X and/or Y each represent straight-chain or branched acyl having up to 4 carbon atoms which is optionally substituted by halogen,

25

or

represent straight-chain or branched acyloxy having up to 4 carbon atoms,

or

represent phenylthio,

and/or

represent radicals of the formulae $-\text{SO}_3\text{H}$ or $\text{S}(\text{O})_c\text{R}^6$,

in which

5 c represents a number 1 or 2,

10 R^6 represents straight-chain or branched alkyl having 1 to 8 carbon atoms, phenyl or a 5- to 6-membered heterocycle having up to 2 heteroatoms from the group consisting of S, N and/or O, where the ring systems may optionally be substituted by halogen or by straight-chain or branched alkyl or alkoxy having in each case up to 3 carbon atoms,

15 and/or the optionally remaining radicals R^1 , X and/or Y each represent a radical of the formula $\text{PO}(\text{OR}^7)(\text{OR}^8)$,

in which

20 R^7 and R^8 are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 4 carbon atoms or cyclopropyl, cyclopentyl, phenyl or benzyl,

25 and/or the optionally remaining radicals R^1 , X and/or Y each represent oxycycloalkyl having 3 to 6 carbon atoms or represent radicals of the formulae $-\text{NH}-\text{C}(=\text{NH})\text{NH}_2$, $-\text{CON}=\text{C}(\text{NH}_2)_2$ or $-\text{C}=\text{NH}(\text{NH}_2)$, $(\text{CO})_d\text{NR}^9\text{R}^{10}$ or $-\text{NHCONR}^9\text{R}^{10}$,

in which


30 d represents a number 0 or 1,

R^9 and R^{10} are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 4 carbon atoms or cyclopropyl, cyclopentyl, cyclohexyl, phenyl or a 3- to 6-membered ring having up to 3 heteroatoms from the group consisting of N, O, S, which may also be attached via N,

where the abovementioned radicals may optionally be substituted phenyl, cyclopropyl, cyclopentyl, hydroxyl, amino or straight-chain or branched alkoxy, acyl or alkoxycarbonyl having in each case up to 4 carbon atoms,

and in the case that $d = 0$,

R^9 and R^{10} may also represent straight-chain, branched or cyclic acyl having up to 6 carbon atoms, straight-chain or branched hydroxyalkyl having up to 4 carbon atoms, straight-chain or branched alkoxyalkyl having a total of up to 10 carbon atoms, straight-chain or branched alkoxycarbonyl or acyloxyalkyl having in each case up to 4 carbon atoms or a radical of the formula $-SO_2R^{11}$ or a radical of the formulae











in which

R^{11} represents straight-chain or branched alkyl having up to 3 carbon atoms,

and/or

R^9 and R^{10} represent radicals of the formulae





in which

R^{12} , R^{13} and R^{15} to R^{28} are identical or different and each represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

e represents a number 0, 1 or 2,

and

R^{14} represents straight-chain or branched alkyl having up to 6 carbon atoms or cycloalkyl having 3 to 8 carbon atoms,

R^9 and R^{10} are identical or different and each represents hydrogen; alkyl having up to 4 carbon atoms which is optionally substituted by hydroxyl or alkoxy having up to 7 carbon atoms, phenyl which is optionally substituted by halogen,

R^2 and R^3 , together with the double bond, form a fused phenyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl ring,

which are optionally substituted up to 2 times by identical or different substituents from the group consisting of formyl, carboxyl, hydroxyl, mercaptyl, straight-chain or branched acyl, alkylthio or alkoxy carbonyl having in each case up to 5 carbon atoms, nitro, cyano, azido, fluorine,

chlorine, bromine or straight-chain or branched alkyl or alkoxy having in each case up to 5 carbon atoms which for its part may be substituted by hydroxyl, amino, carboxyl, straight-chain or branched acyl, alkoxy or alkoxycarbonyl having in each case up to 4 carbon atoms,

5

and/or

the abovementioned heterocyclic rings or phenyl are optionally substituted by a group of the formula $-NR^{29}R^{30}$,

10

in which

R^{29} and R^{30} are identical or different each and represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms

15

or

R^{29} represents hydrogen

20

and

R^{30} represents formyl

25

and/or the abovementioned fused phenyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl rings are optionally substituted by phenyl which for its part may be substituted by fluorine, chlorine, bromine or by straight-chain or branched alkyl or alkoxy having in each case up to 4 carbon atoms,

30

A represents thienyl, tetrahydropyranyl, tetrahydrofuranyl, phenyl, morpholinyl, pyrimidyl, pyrazinyl, pyridazinyl or pyridyl which are optionally substituted up to 2 times by identical or different substituents from the group consisting of hydroxyl, formyl, carboxyl, straight-chain or branched acyl, alkylthio, alkyloxyacyl, alkoxy or alkoxycarbonyl having in each case up to 4 carbon atoms, fluorine, chlorine or bromine,

and their isomeric forms and salts.

Particular preference is given to compounds of the general formula (I) according to the invention,

in which

at least one of the substituents R¹, X and Y represents cyclopropyl which is optionally substituted by hydroxyl or fluoromethyl, or represents cyclobutyl, cyclopentenyl, cyclopentyl or cyclohexyl,

and where the optionally remaining radicals R¹, X and/or Y are identical or different and each represents hydrogen, hydroxyl, halogen or azido,

and/or

represent a 3- to 6-membered ring which may be saturated, unsaturated and/or partially unsaturated and may contain 1 to 3 heteroatoms from the group consisting of N, O, S, SO, SO₂, which may optionally contain a carbonyl group as ring member, which may also be attached via N and which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched alkyl having up to 4 carbon atoms which is optionally substituted by hydroxyl, amino, halogen, carboxyl,

straight-chain or branched acyl, alkoxy, alkoxycarbonyl or acylamino having in each case up to 4 carbon atoms,

and/or

represent straight-chain or branched acyl having up to 4 carbon atoms which is optionally substituted by halogen,

or

represent straight-chain or branched acyloxy having up to 4 carbon atoms,

and/or

represent radicals of the formulae $-SO_3H$ or $S(O)_cR^6$,

in which

c represents a number 1 or 2,

R^6 represents straight-chain or branched alkyl having 1 to 8 carbon atoms, phenyl or a 5- to 6-membered heterocycle having up to 2 heteroatoms from the group consisting of S, N and O, where the ring systems may optionally be substituted by halogen or by straight-chain or branched alkyl or alkoxy having in each case up to 3 carbon atoms,

and/or

represent a radical of the formula $PO(OR^7)(OR^8)$,

in which

R^7 and R^8 are identical or different and each represents hydrogen,
straight-chain or branched alkyl having up to 4 carbon atoms
or cyclopropyl, cyclopentyl, phenyl or benzyl,

and/or

represent oxycycloalkyl having 3 to 6 carbon atoms or represent radicals of
the formulae $-\text{CON}=\text{C}(\text{NH}_2)_2$ or $-\text{C}=\text{NH}(\text{NH}_2)$ or $(\text{CO})_d\text{NR}^9\text{R}^{10}$ or
 $\text{NHCONR}^{12}\text{R}^{13}$,


in which

d represents a number 0 or 1,

R^9 and R^{10} are identical or different and each represent hydrogen,
straight-chain or branched alkyl having up to 4 carbon atoms
or cyclopropyl, cyclopentyl, cyclohexyl or phenyl,

and in the case where $d = 0$

R^9 and R^{10} also represent straight-chain, branched or cyclic acyl
having up to 5 carbon atoms, straight-chain or branched
hydroxyalkyl having up to 3 carbon atoms, straight-chain or
branched alkoxyalkyl having a total of up to 8 carbon atoms,
straight-chain or branched alkoxy carbonyl or acyloxyalkyl
having in each case up to 3 carbon atoms or a radical of the
formula $-\text{SO}_2\text{R}^{11}$ or a radical of the formulae








in which

R^{11} represents straight-chain or branched alkyl having up to 4 carbon atoms

and/or

R^9 and R^{10} represent radicals of the formulae





in which

R^{12} , R^{13} and R^{15} to R^{28} are identical or different and each represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

e represents a number 0, 1 or 2

and

R^{14} represents straight-chain or branched alkyl having up to 6 carbon atoms or cycloalkyl having 3 to 8 carbon atoms,


R^9 and R^{10} are identical or different and each represents hydrogen; alkyl having up to 4 carbon atoms which is optionally substituted by hydroxyl or alkoxy having up to 7 carbon atoms, phenyl which is optionally substituted by halogen,

R^2 and R^3 form, together with the double bond, a phenyl, pyridyl or pyrimidinyl ring,

A represents phenyl or pyrimidyl, each of which is optionally substituted by fluorine, chlorine or bromine,

and their isomeric forms and salts.

In the compounds of the formula (I) according to the invention described above, the radicals R^1 , X and Y are, according to a preferred embodiment, attached to the positions of the pyrimidine ring shown:



R^1 then preferably represents one of the abovementioned optionally substituted cycloalkyl or cycloalkenyl radicals, such as, for example, cyclopropyl, cyclobutyl, cyclopentenyl, cyclopentyl, cyclohexyl, 1-hydroxycyclopropyl and 1-(fluoromethyl)cyclopropyl; amongst these, cyclopropyl is particularly preferred.

X then represents, for example, an NH_2 group and

Y represents hydrogen or an NH_2 group.

5

In the compounds of the formula (I) according to the invention described above, R^2 and R^3 represent, according to a preferred embodiment and together with the double bond, a fused phenyl ring or a fused pyridyl ring, resulting in an indazole or a pyrazolo[3,4]-pyridine skeleton. Particular preference is given to a pyrazolo[3,4]-pyridine skeleton.

10


In the compounds of the formula (I) according to the invention described above, according to a preferred embodiment, the radical A represents a halogen-, in particular fluorine-, substituted phenyl ring. The substituent is preferably in the ortho-position to the methylene bridge.

15

In addition, processes for preparing compounds according to the invention of the general formula (I) have been found, characterized in that, depending on the various meanings of the heterocycles listed above under R^2 and R^3 ,

20

[A] compounds of the general formula (II)



25


in which

R^1 , X and Y are each as defined above

and

D represents radicals of the formulae

5



in which

10

R^{35} represents $\text{C}_1\text{-C}_4$ -alkyl

are converted, by reaction with compounds of the general formula (III)




15


in which

A is as defined above

20



(IV) and



(IVa),

in which


A, X, Y and R¹ are each as defined above,

and, in the case of the compounds of the general formula (IVa), are subsequently cyclized with carboxylic acids, nitriles, formamides or guanidium salts,

and, in the case of the compounds of the general formula (IV), are cyclized with 1,3-dicarbonyl derivatives, their salts, tautomers, enol ethers or enamines, in the presence of acids and, if appropriate, under microwave irradiation,

or


[B] in the case that R² and R³ together form a pyrazine ring, compounds of the general formula (IV) are initially converted by nitrosation into the compounds of the general formula (V)



in which

A, X, Y and R¹ are each as defined above,

in a second step, the compounds of the general formula (VI)



in which


A, X, Y and R¹ are each as defined above

are prepared by a reduction,

and these are subsequently cyclized with 1,2-dicarbonyl compounds, preferably aqueous glyoxal solution,

or

[C] compounds of the general formula (VII)



in which

A^1 , R^2 and R^3 are each as defined above

and

5 L represents a radical of the formula $-\text{SnR}^{36}\text{R}^{37}\text{R}^{38}$, ZnR^{39} , iodine, bromine or triflate,


in which

10 R^{36} , R^{37} and R^{38} are identical or different and each represent straight-chain or branched alkyl having up to 4 carbon atoms

and

15 R^{39} represents halogen

are reacted with compounds of the general formula (VIII)



in which

25 X, Y and R^1 are each as defined above

and

in the case that $L = \text{SnR}^{36}\text{R}^{37}\text{R}^{38}$ or ZnR^{39} ,

T represents triflate or represents halogen, preferably bromine

and,

in the case that $L = \text{iodine, bromine or triflate,}$

T represents a radical of the formula $\text{SnR}^{36}\text{R}^{37}\text{R}^{38}$, ZnR^{39} or $\text{BR}^{40}\text{R}^{41}$,

in which


R^{36} , R^{37} , R^{38} and R^{39} have the meanings of R^{36} , R^{37} , R^{38} and R^{39} given above and are identical to or different from them,

R^{40} and R^{41} are identical or different and each represent hydroxyl, aryloxy having 6 to 10 carbon atoms or straight-chain or branched alkyl or alkoxy having in each case up to 5 carbon atoms, or together form a 5- or 6-membered carbocyclic ring

in a palladium-catalysed reaction in inert solvents, if appropriate in the presence of a base,

or


[D] amidines of the general formula (IX)



in which

5 A, R² and R³ are each as defined above


are reacted, for example, with compounds of the general formula (X), (Xa), (Xb) or (Xc)




(X)



(Xa)



(Xb)



(Xc)

in which

R^{1'} represents the optionally substituted cycloalkyl radical listed above under R¹;

Alk represents straight-chain or branched alkyl having up to 8 carbon atoms, preferably up to four carbon atoms;

and


Z represents an NH_2 group, a monoalkylamino group having up to 7 carbon atoms, a dialkylamino group having up to 7 carbon atoms, a piperidinyl or morpholinyl radical which is attached via the nitrogen, hydroxyl, alkoxy having up to 7 carbon atoms, acyloxy having up to 7 carbon atoms or aroyloxy having 6 to 10 carbon atoms,

and, if appropriate, the substituents listed under X, Y, R^1 , R^2 , R^3 and/or A are modified or introduced by customary methods, preferably by acylation and derivatization of free amino groups, chlorination, catalytic hydrogenation, reduction, oxidation, removal of protective groups and/or nucleophilic substitution.

The heterocycles listed under R^2 and R^3 can also be introduced by reacting the appropriately substituted compounds of the general formula (II) according to other known heterocyclic syntheses.

The process [D] according to the invention can be illustrated by the following embodiment:

[D]



Suitable solvents for the individual steps of process [A] and [B] are inert organic solvents which do not change under the reaction conditions. These include ethers, such as diethyl ether or tetrahydrofuran, DME, dioxane, alcohols, such as methanol and ethanol, halogenated hydrocarbons, such as dichloromethane, trichloromethane, carbon tetrachloride, 1,2-dichloroethane, trichloroethane, tetrachloroethane, 1,2-dichloroethylene or trichloroethylene, hydrocarbons, such as benzene, xylene, toluene, hexane, cyclohexane, or mineral oil fractions, nitromethane, dimethylformamide, acetone, acetonitrile or hexamethylphosphoric triamide. It is also possible to use mixtures of the solvents. Particular preference is given to tetrahydrofuran, dimethylformamide, toluene, dioxane or dimethoxyethane.

Bases which are suitable for use in the processes according to the invention are, in general, inorganic or organic bases. These preferably include alkali metal hydroxides, such as, for example, sodium hydroxide or potassium hydroxide, alkaline earth metal hydroxides, such as, for example, barium hydroxide, alkali metal carbonates, such as sodium carbonate or potassium carbonate, alkaline earth metal carbonates, such as

calcium carbonate, or alkali metal or alkaline earth metal alkoxides, such as sodium methoxide or potassium methoxide, sodium ethoxide or potassium ethoxide or potassium tert-butoxide, or organic amines (trialkyl(C_1-C_6)-amines), such as triethylamine, or heterocycles, such as 1,4-diazabicyclo[2.2.2]octane (DABCO),
 5 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), pyridine, diaminopyridine, methylpiperidine or morpholine. It is also possible to use alkali metals, such as sodium, and their hydrides, such as sodium hydride, as bases. Preference is given to sodium carbonate and potassium carbonate, triethylamine and sodium hydride.

10 When reacting the compounds of the formula (II) with the compounds of the formula (III), the base is employed in an amount of from 1 mol to 5 mol, preferably from 1 mol to 3 mol, based on 1 mol of the compound of the general formula (II).

The reaction of the compounds of the formula (II) with the compounds of the formula
 15 (III) is generally carried out in a temperature range of from 0°C to 150°C, preferably from +20°C to +110°C.

This reaction can be carried out at atmospheric pressure or at elevated or reduced pressure (for example from 0.5 to 5 bar). In general, the reaction is carried out at
 20 atmospheric pressure.

Suitable acids for the cyclization reactions which may have to be carried out in the processes according to the invention are, in general, protic acids. These preferably include inorganic acids, such as, for example, hydrochloric acid or sulphuric acid, or
 25 organic carboxylic acids having 1-6 C atoms which are optionally substituted by fluorine, chlorine and/or bromine, such as, for example, acetic acid, trifluoroacetic acid, trichloroacetic acid or propionic acid, or sulphonic acids having C_1-C_4 -alkyl radicals or aryl radicals, such as, for example, methanesulphonic acid, ethanesulphonic acid, benzenesulphonic acid or toluenesulphonic acid.

5 The catalytic hydrogenation reactions which may have to be carried out in the processes according to the invention can generally be carried out with hydrogen in water or in inert organic solvents, such as alcohols, ethers or halogenated hydrocarbons, or mixtures thereof, using catalysts such as raney nickel, palladium, palladium on animal charcoal or platinum, or using hydrides or boranes in inert solvents, if appropriate in the presence of a catalyst.

10 The chlorination reactions which may have to be carried out in the processes according to the invention are generally carried out using the customary chlorinating agents, such as, for example, PCl_3 , PCl_5 , POCl_3 or elemental chlorine. In the context of the invention, preference is given to POCl_3 .

15 The acylations and derivatizations of free amino groups which may have to be carried out in the processes according to the invention can be carried out by customary methods which are known to the person skilled in the art. It is possible, for example, to convert appropriate free amino groups by reaction with an acyl halide, preferably an acyl chloride, or an acetic anhydride, in the presence of a base, such as, for example, sodium hydride, pyridine or dimethylaminopyridine in a solvent such as tetrahydrofuran or dichloromethane into the respective amides, by reaction with an appropriate aldehyde in a solvent such as ethanol or acetonitrile into the respective
20 N,O-hemiacetals or N,O-acetals by reaction with a sulphonyl halide, preferably a sulphonyl chloride, into the respective sulphonamides, by reaction with a chloroformic ester into the respective urethanes or by reaction with an isocyanate in a solvent such as dichloromethane into the respective urea derivatives.

25 The nucleophilic substitutions and Vilsmeier reactions which may have to be carried out in the processes according to the invention are carried out by customary methods known to the person skilled in the art.

The nitrosation of the compounds of the formula (IV) to the compounds of the formula (V), which constitutes the first step of the process [B], can be carried out in accordance with the procedure of P.G. Baraldi et al., Synthesis 1984, 148.

5 The reductions which may have to be carried out in the processes according to the invention are generally carried out using reducing agents, preferably those which are suitable for reducing carbonyl to hydroxyl compounds. Particularly suitable here is the reduction with metal hydrides or complex metal hydrides in inert solvents, if appropriate in the presence of a trialkylborane. Preference is given to reduction with complex metal hydrides, such as, for example, lithium borohydride, sodium borohydride, potassium borohydride, zinc borohydride, lithium trialkyl borohydride, diisobutylaluminium hydride or lithium aluminium hydride. The reduction is very particularly preferably carried out using diisobutylaluminium hydride and sodium borohydride.

10 Here, the reducing agent is generally employed in an amount of from 1 mol to 6 mol, preferably from 1 mol to 4 mol, based on 1 mol of the compounds to be reduced.

15 The reductions which may have to be carried out in the processes according to the invention are generally carried out in a temperature range of from -78°C to $+50^{\circ}\text{C}$, preferably from -78°C to 0°C in the case of DIBAH and 0°C to room temperature in the case of NaBH_4 .

20 The reductions which may have to be carried out in the processes according to the invention are generally carried out at atmospheric pressure. However, it is also possible to operate under elevated or reduced pressure.

25 The compounds of the general formulae (II) and (III) are known per se, or they can be prepared by customary methods (cf.: J. Hromatha et al., Monatsh. Chem. 1976, 107, 233).

09744703-032601

Some of the compounds of the general formulae (IV), (IVa), (V) and (VI) are known, and they can be prepared as described above.

Suitable solvents for the process [C] are inert organic solvents which do not change under the reaction conditions. These include ethers, such as diethyl ether or tetrahydrofuran, DME, dioxane, halogenated hydrocarbons, such as dichloromethane, trichloromethane, carbon tetrachloride, 1,2-dichloroethane, trichloroethane, tetrachloroethane, 1,2-dichloroethylene or trichloroethylene, hydrocarbons, such as benzene, xylene, toluene, hexane, cyclohexane, or mineral oil fractions, nitromethane, dimethylformamide, acetone, acetonitrile or hexamethylphosphoric triamide. It is also possible to use mixtures of the solvents. Particular preference is given to tetrahydrofuran, dimethylformamide, toluene, dioxane or dimethoxyethane.

The reaction of the compounds of the formula (VII) with the compounds of the formula (VIII) is generally carried out in a temperature range of from 0°C to 150°C, preferably from +20°C to +110°C.

This reaction can be carried out at atmospheric pressure or at elevated or reduced pressure (for example from 0.5 to 5 bar). In general, the reaction is carried out at atmospheric pressure.

Suitable palladium compounds in the context of the present invention are, in general, $\text{PdCl}_2(\text{P}(\text{C}_6\text{H}_5)_3)_2$, palladium-bis-dibenzylideneacetone ($\text{Pd}(\text{dba})_2$), [1,1'-bis-(diphenylphosphino)ferrocene]-palladium(II) chloride ($\text{Pd}(\text{dppf})\text{Cl}_2$) or $\text{Pd}(\text{P}(\text{C}_6\text{H}_5)_3)_4$. Preference is given to $\text{Pd}(\text{P}(\text{C}_6\text{H}_5)_3)_4$.

The compounds of the general formula (VII) are known per se, or they can be prepared by customary methods (cf., for example K. Kirschke in: Houben-Weyl, Methoden der organischen Chemie, Thieme-Verlag Stuttgart, 4th Ed., volume E8b, part 2, 399-763; in particular with respect to pyrazolopyridines: C.R. Hardy in A.R. Katritzky (Ed.), Adv. Het. Chem. 1984, 36, 343-409; in particular with respect to pyrazolopyrimidines:

M.H. Elgnadi et al., Adv. Het. Chem. 1987, 41, 319-376). The preparation of the corresponding halogenopyrazolo[3,4-b]pyrimidines and organotin pyrazolo[3,4-b]pyrimidines of the formula (VII) is described in WO 98/23619 and can also be carried out analogously for the corresponding triflate and organotin compounds of the formula (VII).


The compounds of the general formula (VIII) are known and can be prepared by customary methods (cf., for example, M.G. Hoffmann et al. in: Houben-Weyl, Methoden der organischen Chemie, 4th ed., volume E9b, part 1, pp. 1-249; A. Weissenberger et al., The Chemistry of heterocyclic compounds – pyrimidines, 1962, 16; ibid 1970, 16, suppl. 1, ibid 1985, 16, suppl. 2; ibid 1994, 52).

The process [D] is carried out in a temperature range of from 80°C to 120°C, preferably at from 100°C to 110°C, or under reflux.

Suitable solvents are, for example, the reagents of the general formula (X), (Xa), (Xb) or (Xc). However, the reaction can also be carried out in other suitable solvents, such as, for example, toluene, methanol or dichloromethane. Low-boiling solvents, such as, for example, dichloromethane, can be distilled off during the course of the reaction.

The process [D] can be carried out at atmospheric pressure or at elevated or reduced pressure (for example from 0.5 to 5 bar). In general, the process is carried out at atmospheric pressure.

The amidines of the general formula (IX) are novel and are therefore a further subject of the invention. They can be prepared by reacting the compounds of the general formula (XI)




in which

A, R² and R³ are each as defined above

5

initially in ethers with trifluoroacetic anhydride (TFAA) and in the presence of bases to give the compound of the general formula (XII)




in which

A, R² and R³ are each as defined above,

10


15 subsequently preparing the compounds of the general formula (XIII)



in which

A, R² and R³ are each as defined above

using sodium methoxide, in a next step converting these compounds by reaction with NH₄Cl and glacial acetic acid in alcohols into the corresponding amidine HCl salt of the general formula (XIV)



in which

A, R² and R³ are each as defined above

and, in a last step, reacting with bases, preferably sodium carbonate, or alkali metal alkoxide, such as sodium ethoxide.

Suitable solvents for reacting the compounds of the general formula (XI) to give the compounds of the formula (XII) are ethers, such as diethyl ether or tetrahydrofuran, dimethylformamide and dioxane; preference is given to tetrahydrofuran.

Suitable for use as bases here are organic amines (trialkyl(C₁-C₆)-amines) such as triethylamine, or heterocycles, such as 1,4-diazabicyclo[2.2.2]octane (DABCO), 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), pyridine, dimethylaminopyridine, methylpiperidine or morpholine. Preference is given to pyridine.

The reaction is carried out in a temperature range of from 0°C to 40°C, preferably at room temperature.

The reaction can be carried out at atmospheric pressure or at elevated or reduced pressure (for example from 0.5 to 5 bar). In general, the reaction is carried out at atmospheric pressure.

5

The amide (XI) can be prepared, for example, by hydrolysing an appropriate ester as starting material with a base to give the acid, converting the acid into the acyl chloride by customary methods, for example using SOCl_2 or POCl_3 , followed by reaction with ammonia.

10

The elimination of water from the amide (XI) to given the nitrile (XII) can be carried out with any customary dehydrating agent. Preference according to the invention is given to trifluoroacetic anhydride (TFAA).

15

The nitrile of the formula (XII) can be converted into the iminoether of the formula (XIII) both under acidic conditions, such as, for example, with HCl /alcohol mixtures, and under basic conditions, such as, for example, with methanol/sodium methoxide. It is generally carried out at from 0°C to 40°C , for example at room temperature.

20

Suitable solvents for converting the compounds of the general formula (XIII) into the compounds of the formula (XIV) are alcohols, such as methanol or ethanol. Preference is given to methanol.

25

The reaction is carried out in a temperature range of from 0°C to 40°C , preferably at room temperature.

30

The reaction can be carried out under atmospheric pressure or under elevated or reduced pressure (for example from 0.5 to 5 bar). In general, the reaction is carried out at atmospheric pressure.


Suitable bases for liberating the compounds of the general formula (IX) from compounds of the general formula (XIV) are inorganic or organic bases. These include, for example, alkali metal hydroxides, such as sodium hydroxide or potassium hydroxide, alkaline earth metal hydroxides, such as barium hydroxide, alkali metal carbonates, such as sodium carbonate or potassium carbonate, alkaline earth metal carbonates, such as calcium carbonate, and alkali metal alkoxides, such as sodium methoxide. Preference is given to sodium carbonate and sodium methoxide.

The pyrimidine ring is prepared by customary methods (cf., for example, M.G. Hoffmann et al. in: Houben-Weyl, Methoden der organischen Chemie, 4th ed., volume E9b, part 1, pp. 1-249; A. Weissenberger et al., The Chemistry of heterocyclic compounds - pyrimidines, 1962, 16; ibid 1970, 16, suppl. 1, ibid 1985, 16, suppl. 2; ibid 1994, 52).


Here, the iminoethers of the formula (XIII) can be used as starting materials and be reacted, for example, with a suitable enamine. However, it is also possible to convert the iminoether first, using ammonia or its salts, into a corresponding amidine and to react this either as the free base (IX) or as a salt (XIV) with enamines, acetals, enol ethers, aldehydes, enolates, malononitrile esters or malonodinitriles.

The enamines which may have to be used in this reaction can be prepared, for example, from C-H-acidic compounds, such as acetonitrile derivatives, according to known methods, by reaction with dimethylformamide derivatives, such as, for example, bis(dimethylamino)-tert-butoxymethane, dialkoxy-dialkylamino-methanes.


The compounds of the general formula (XI) are novel and are therefore a further subject of the invention. They can be prepared by converting the compounds of the general formula (XV)



with the compounds of the general formula (XVI)




in ethers, preferably dioxane, and trifluoroacetic acid into the compounds of the general formula (XVII)



10

subsequently preparing, by reaction with the compounds of the general formula (XVIII)




in which

Z is as defined above, in particular $-\text{N}(\text{CH}_3)_2$

20

in inert solvents, preferably dioxane, the compounds of the general formula (XIX)




followed in a last step by treatment with ammonia in methanol.


- 5 Instead of the sodium salt of the enolate (XV), it is also possible to employ enol ethers, ketones or enamines.

If appropriate, the reaction of the compounds of the general formulae (XV) and (XVI) to give (XVII) can also be carried out via intermediates of the formulae (A) and (B), which are likewise novel,

10



A





B

at room temperature. These represent, therefore, a further subject of the invention.

15

The compounds of the general formula (X) are novel and can be prepared, for example, by reacting the compounds of the formula (XX) or (XXa)



20

in which

Alk and Alk⁺ are identical or different and each represent straight-chain or branched alkyl having up to 5 carbon atoms

5 with compounds of the formula (XXI)



in which

10 R^1 represents the cycloalkyl radical listed above under R^1 .

The compounds of the general formulae (XX), (XXa) and (XXI) are known, or they can be prepared by customary methods.


15

Some of the compounds of the general formulae (XII), (XIII), (XIV), (XV), (XVII), (XVIII) and (XIX) are novel, and they can be prepared as described above.

20

The pyrimidine radical can also be synthesized with the aid of the reagent of the formula (Xa) which is accessible, for example, as follows:

Compounds of the general formula




25

in which

$R^{1'}$ is as defined above and Alk represents an alkyl radical having up to 4 carbon atoms

are converted, by using ammonia in suitable solvents, preferably alcohols such as methanol, at temperatures from 0°C to 40°C, preferably at room temperature, into compounds of the general formula (XXIII)




in which $R^{1'}$ is as defined above,

and these are subsequently reacted by customary methods with dehydrating agents, such as, for example, Burgess reagent, POCl_3 , P_2O_5 , SOCl_2 , trifluoroacetic anhydride/pyridine.


If Burgess reagent is used, the reaction is preferably carried out in inert solvents, such as ethers or chlorinated hydrocarbons. Examples which may be mentioned are dichloromethane and tetrahydrofuran. Preference is given to using a 1:2 mixture of the abovementioned solvents. The reaction is carried out at temperatures from 0°C to 40°C, preferably at room temperature.

The compounds of the formula (XXII) are known and/or obtainable in a simple manner known to the person skilled in the art.

Some of the compounds of the formula (X) undergo keto-enol-tautomerism, for example:



If R^2 and R^3 together with the double bond form a phenyl ring, the corresponding 3-cyano-indazoles are reacted with compounds of the general formula (XXIV)



in which

A is as defined above


10

and

Hal represents Cl, Br or I, preferably Br,

15

in inert solvents, preferably with tetrahydrofuran in the presence of a base, preferably sodium hydride, to give the compounds of the general formula (XXV)



20

in which

A is as defined above

which are subsequently treated with ammonium chloride and sodium methoxide as described above.

25

The compounds of the general formula (XXIV) are known, or they can be prepared by customary methods.

5 The compounds of the general formula (XXV) are novel and can be prepared as described above.

The compounds of the general formula (I) according to the invention have an unforeseeable, valuable spectrum of pharmacological action.

10 The compounds of the general formula (I) according to the invention lead to a vasorelaxation, inhibition of platelet aggregation and to a fall in blood pressure and also to an increase in coronary blood flow. These actions are mediated via a direct stimulation of soluble guanylate cyclase and an intracellular cGMP increase. Additionally, the compounds of the general formula (I) according to the invention
15 enhance the action of substances which raise the cGMP level, such as, for example, EDRF (endothelium-derived relaxing factor), NO donors, protoporphyrin IX, arachidonic acid or phenylhydrazine derivatives.

20 They can therefore be employed in medicaments for the treatment of cardiovascular disorders, such as, for example, for the treatment of high blood pressure and cardiac insufficiency, stable and unstable angina pectoris, peripheral and cardiac vascular disorders, arrhythmias, for the treatment of thromboembolic disorders and ischaemias such as myocardial infarct, stroke, transitory and ischaemic attacks, peripheral circulatory disorders, prevention of restenoses such as after thrombolysis therapy,
25 percutaneous translumino angioplastie (PTA), percutaneous transluminalo coronary angioplasty (PTCA), bypass and also for the treatment of arterioscleroses, asthmatic disorders and disorders of the urogenital system such as, for example, prostate hypertrophy, erectile dysfunction, female sexual dysfunction and incontinence.

30 The compounds of the general formula (I) described in the present invention are also active compounds for controlling disorders in the central nervous system which are

characterized by disturbances of the NO/cGMP system. In particular, they are suitable for eliminating cognitive deficits, for improving learning and memory performance and for treating Alzheimer's disease. They are also suitable for the treatment of disorders of the central nervous system such as states of anxiety, tension and depression, sleeping disorders and sexual dysfunction caused by the central nervous system, and for regulating pathological eating disorders or disorders associated with the use of stimulants and drugs.

Furthermore, the active compounds are also suitable for regulating cerebral circulation, and they are therefore effective agents for controlling migraines.

They are also suitable for the prophylaxis and control of the sequelae of cerebral infarcts (Apoplexia cerebri) such as stroke, cerebral ischaemia and skull-brain trauma. The compounds of the general formula (I) according to the invention can also be employed for controlling pain.

Additionally, the compounds according to the invention have antiinflammatory action and can therefore be employed as antiinflammatories.

The invention moreover includes the combination of the compounds of the general formula (I) according to the invention with organic nitrates and NO donors.

Organic nitrates and NO donors in the context of the invention are, in general, substances which display their therapeutic action by the release of NO or NO species. Sodium nitroprusside, glycerol trinitrate, isosorbide dinitrate, isosorbide mononitrate, molsidomine and SIN-1 are preferred.

The invention additionally includes the combination with compounds which inhibit the degradation of cyclic guanosine monophosphate (cGMP). These are, in particular, inhibitors of phosphodiesterases 1, 2 and 5; nomenclature according to Beavo and Reifsnnyder (1990) TIPS 11 p. 150 to 155. The action of the compound according to the

invention is potentiated and the desired pharmacological effect is increased by these inhibitors.

To determine the cardiovascular action, the following investigations were carried out:

- 5 In in vitro investigations on the isolated enzyme and on cells of vascular origin, the effect on guanylate cyclase-dependent cGMP formation was tested with and without NO donor. The anti-aggregatory properties were shown on human platelets stimulated with collagen. The vasorelaxant action was determined in rabbit aortal rings precontracted with phenylephrine. The hypotensive action was investigated in
10 anaesthetized and awake rats.

Stimulation of recombinant soluble guanylate cyclase in vitro

The investigations on the stimulation of recombinant soluble guanylate cyclase by the compounds according to the invention with and without NO donor were carried out using the method which is described in detail in the following literature reference: M. Hoenicka, E.M. Becker, H. Apeler, T. Sirichoke, H. Schroeder, R. Gerzer and J.-P. Stasch: Purified soluble guanylate cyclase expressed in a baculovirus/Sf9 system: stimulation by YC-1, nitric oxide, and carbon oxide. J. Mol. Med. 77: 14-23 (1999). The results are shown in Fig. 1.

Stimulation of soluble guanylate cyclase in primary endothelial cells

Primary endothelial cells were isolated from pig aortas by treatment with collagenase soln. The cells were then cultured in culture medium at 37°C/5% CO₂ until confluence was reached. For the investigations, the cells were passaged, inoculated into 24-well cell culture plates and subcultured until reaching confluence (~ 2 x 10⁵ cells/well). For the stimulation of endothelial guanylate cyclase, the culture medium was aspirated and the cells were washed once with Ringer solution. After removing the Ringer solution, the cells were incubated for 10 minutes, at 37°C/5% CO₂ in stimulation buffer. Following this, the test substances (final concentration 1 µM) were added to the cells by pipette. After a further 10 minutes the buffer solution was aspirated and cold buffer at 4°C was added to the cells. The cells were then lysed at -20°C for 16 hours. The supernatants containing the intracellular cGMP were then removed and the cGMP concentration was determined by means of the cGMP-SPA system (Amersham Buchler, Brunswick). The results are shown in Table 1.

Table 1:

Ex. No.	Concentration (μM)	0	0.1	0.3	1.0	3.0	10
1	cGMP (pmol/well)	1.4	2.2	4.0	6.9	8.5	14.6

5 Vasorelaxant action in vitro

Rabbits are anaesthetized by a blow to the neck and exanguinated. The aorta is removed, freed from adhering tissue, divided into 1.5 mm wide rings and individually transferred under a pretension into 5 ml organ baths containing a warm, carbogen-aerated Krebs-Henseleit solution at 37°C of the following composition (mM): NaCl: 119; KCl: 4.8; $\text{CaCl}_2 \times 2 \text{ H}_2\text{O}$: 1; $\text{MgSO}_4 \times 7 \text{ H}_2\text{O}$: 1.4; KH_2PO_4 : 1.2; NaHCO_3 : 25; glucose: 10. The contractility is detected using Satham UC2 cells, amplified and digitalized by means of A/D converters (DAS-1802 HC, Keithley Instruments Munich), and recorded in parallel on linear recorders. To produce a contraction, phenylephrin is added to the bath cumulatively in increasing concentration. After several control cycles, the substance to be investigated is investigated in each further passage in increasing dosage in each case, and the height of the contraction is compared with the height of the contraction achieved in the last preliminary passage. From this, the concentration which is necessary in order to reduce the height of the control value by 50% (IC_{50}) is calculated. The standard administration volume is 5 μl , and the proportion of DMSO in the bath solution corresponds to 0.1%. The results are shown in Table 2.

Table 2

Example No.	IC_{50} [μM]
1	0.2 μM

Blood pressure measurements in anaesthetized rats

Male Wistar rats having a bodyweight of 300-350 g are anaesthetized with Thiopental (100 mg/kg i.p.). After tracheotomy, a catheter is inserted into the femoral artery for blood pressure measurement. The substances to be tested are administered orally in Transcutol, Cremophor EL, H₂O (10%/20%/70%) in a volume of 1 ml/kg. The results are listed in Table 3 below.

Table 3

Ex. No.	Dose (mg/kg p.o.)	Max. reduction in blood pressure (mm Hg)	Time (min)
1	1	23	20
1	3	37	40

Effect on the average blood pressure of awake, spontaneously hypertensive rats

Continuous measurements of blood pressure over 24 hours were carried out on spontaneously hypertensive female rats (MOL:SPRD) having a bodyweight of 200-250 g which were allowed to move around freely. To this end, pressure monitors (Data Sciences Inc., St. Paul, MN, USA) were chronically implanted into the descending abdominal aorta of the animals, below the kidney artery, and the attached transmitter was fixed in the abdominal cavity.

The animals were kept individually in type III cages, which were positioned on the individual receiver stations, and adapted to a 12-hour day/night rhythm. Water and feed was available freely.

To collect the data, the blood pressure of each rat was registered every 5 minutes for 10 seconds. In each case, the data for a period of 15 minutes were collected and the average value was calculated from these values.

- 5 The compounds to be tested were dissolved in a mixture of Transcutol (10%), Cremophor (20%), H₂O (70%) and administered orally by means of a stomach tube in a volume of 2 ml/kg of bodyweight. The doses tested were between 0.3 and 30 mg/kg of bodyweight. The results are shown in Fig. 2.

10 **Inhibition of platelet aggregation in vitro**

To determine the platelet-aggregation, blood from healthy volunteers of both sexes was used. As an anticoagulant, 9 parts of blood were admixed to one part of 3.8% strength sodium citrate solution. The blood was centrifuged at 900 rpm for 20 min.

- 15 The pH of the platelet-rich plasma obtained was adjusted to pH 6.5 using ACD solution (sodium citrate/citric acid/glucose). The platelets were subsequently centrifuged off and resuspended in buffer and once more centrifuged off. The platelet pellet was suspended in buffer and mixed with an additional 2 mmol/l of CaCl₂.

- 20 To measure the aggregation, aliquots of the platelet suspension were incubated with the substance to be tested at 37°C for 10 min. Aggregation was subsequently induced in an aggregometer by addition of collagen and determined at 37°C using the turbidometric method according to Born (Born, G.V.R., J.Physiol. (London), 168, 178-195, 1963). The results are shown in Table 2 below.

Table 4

Ex. No.	IC ₅₀ (μM)
1	0.003

Measurement of the erection-promoting action of guanylate cyclase stimulators

5

For a complete and lasting erection to occur, the cavernous arteries and the entire cavernous body architecture, which is formed by a network of smooth muscle cells and collagen connective tissue, has to be at maximum dilation so that the corpus cavernosum can fill completely with blood (Anderson K.-E. and Wagner G.,
 10 "Physiology of Penile Erection." *Physiological Reviews* 75, 191-236 (1995);
 Meinhardt W. Kropmann RF, Vermeig P, Lyecclama a Nigelholt and Zwartendijk J.
 "The Influence of Medication on Erectile dysfunction." *Int. J. of Impotence Res.* 9,
 17-26 (1997). Relaxation of smooth muscles is mediated by NO which, in the case of
 15 sexual stimulation, is released by non-adrenergic, non-cholinergic nerve fibres in the
 endothelial cells of the blood vessels of the corpus cavernosum. NO activates
 guanylate cyclase, and the resulting increase in cGMP leads to dilation of the smooth
 muscles of the corpus cavernosum and consequently to an erection. To test the
 efficacy of the substances according to the invention, awake rabbits were used. The
 species rabbit was chosen since neurophysiology, haemodynamic and the control of
 20 contraction and relaxation of the smooth muscles of the corpus cavernosum of rabbit
 and man are quite similar (Meyer MF, Taher H., Krah H., Staubesand J., Becker AJ,
 Kircher M, Mayer B., Jonas U., Forsmann WG., Stief Ch.G. "Intracavenous
 Application of SIN-1 in Rabbit and Man: Functional and Toxological Results."
Annals Urol. 27, 179-182 (1993); Taub HC, Lerner SE, Melman A, Christ GJ
 25 „Relationship between contraction and relaxation in human and rabbit corpus
 cavernosum." *Urology* 42, 698-671, (1993).

Method:

Adult male chinchilla rabbits having a weight of 3-5 kg are, after delivery, adapted for several days in isolation. They have free access to water and can feed for two hours per day. The animals are kept in a 10/14 hour day/night rhythm (light switched on from 8.00 hours onwards). The room temperature is 22 -24 °C.

The animals are weighed directly before the start of the experiment. For intravenous administration, the substances according to the invention were dissolved in a mixture of Transcutol (GATTEFOSSE GmbH) diluted with 20% Cremophor (BASF) and water in a ratio of 3/7. Sodium nitroprusside was dissolved in 0.9% NaCl. The substances were injected at the dosages stated in the table in a volume of 0.5 ml/kg into the auricular vein. For oral administration, the test substances were dissolved in a mixture of glycerol: water: polyethylene glycol 6:10:9.69 and administered at the dosages stated in the table in a volume of 1 ml/kg using the stomach tube.

The effect of guanylate cyclase stimulators is increased by NO donors. This was demonstrated by additionally administering sodium nitroprusside.

The sodium nitroprusside was injected into the auricular vein at a dosage of 0.2 mg/kg simultaneously with the substance according to the invention. If the substance according to the invention was administered orally, the sodium nitroprusside was injected into the auricular vein of these animals 30 min after the oral administration. Corresponding controls were carried out using the solvent and using sodium nitroprusside on its own.

At rest, the penis of the rabbit is not visible in the pubic region and is covered completely by the sheath. The erection is assessed by measuring the length of the protruding penis with a calliper square. Measurements are carried out 5, 10, 15, 30, 45, 60, 120 and 180 min. after the administration of the substance. The effect is

calculated as the product of the length of the penis which is not covered by fur in [mm] and the time for which the erection persists in [min].

- 5 Intravenous injection of sodium nitroprusside causes an erection which lasts for approximately 10 min. (110 [mm × min]).

Table 5: Erection-promoting effect of guanylate cyclase stimulators

(Unit: [mm] × [min])

Substance	Dose i.v.	+SNP i.v.	-SNP i.v.	Dose po	+SNP po	-SNP po
Ex. 1	0.3 mg/kg	140	0	1 mg/kg	184	ng
	1 mg/kg	369	184	3 mg/kg	233	ng
				10 mg/kg	375	ng
				30 mg/kg	804	418
Ex. 25	0.3 mg/kg	85	0	3 mg/kg	168	ng
	1 mg/kg	245	0	10 mg/kg	506	ng

Remarks:

5

10

The substances were administered in the dosages stated. The sodium nitroprusside was in each case administered i.v., at 0.2 mg/kg. If the substance according to the invention was administered i.v., the sodium nitroprusside was administered simultaneously, if the substance according to the invention was administered orally, it was administered 30 min later. Sodium nitroprusside on its own has an effect of 100 [mm]×[min].

15

The present invention includes pharmaceutical preparations which, in addition to non-toxic, inert pharmaceutically acceptable excipients, contain the compounds of the general formula (I) according to the invention, and also processes for the production of these preparations.

20

The active compounds can optionally be present in one or more of the excipients indicated above and also in microencapsulated form.

The therapeutically active compounds of the general formula (I) should be present in the abovementioned pharmaceutical preparations in a concentration from

approximately 0.1 to 99.5, preferably from approximately 0.5 to 95, % by weight of the total mixture.

5 In addition to the compounds of the general formula (I) according to the invention, the abovementioned pharmaceutical preparations can also contain other pharmaceutically active compounds.

10 In general, it has proved advantageous both in human and in veterinary medicine to administer the active compound(s) according to the invention in total amounts from approximately 0.5 to approximately 500, preferably 5 to 100, mg/kg of bodyweight every 24 hours, if appropriate in the form of several individual doses, to achieve the desired results. An individual dose contains the active compound(s) according to the invention preferably in amounts from approximately 1 to approximately 80, in particular 3 to 30, mg/kg of bodyweight.

15 Below, the present invention is illustrated in more detail using non-limiting, preferred examples. Unless indicated otherwise, all amounts given refer to per cent by weight.

Examples**Abbreviations:**

- 5 RT: Room temperature
 EA: Ethyl acetate
 BABA: n-Butyl acetate/n-butanol/glacial acetic acid/phosphate buffer pH 6
 (50:9:25:15; org. phase)

10 **Mobile phases for thin-layer chromatography:**

- T1 E1: toluene/ethyl acetate (1:1)
 T1 EtOH1: toluene-methanol (1:1)
 C1 E1: cyclohexane/ethyl acetate (1:1)
 15 C1 E2: cyclohexane/ethyl acetate (1:2)

Starting materials**Example I: Cyclopropyl-3-oxopropionitrile**


20

- Under argon, 65.4 g (0.583 mol) of potassium tert-butoxide are dissolved in 400 ml of THF. At room temperature, a solution of 21.5 g (0.265 mol) of cyclopropylacetonitrile and 43.2 g (0.58 mol) of ethyl formate in 100 ml of THF is added. The mixture is
 25 stirred at RT for 3 hours. The THF is stripped off using a rotary evaporator, and the residue is partitioned between 200 ml of ice water and 200 ml of ethyl acetate. The organic phase is discounted, whereas the aqueous phase is adjusted to pH = 4 using hydrochloric acid and extracted twice with ethyl acetate. The organic extracts are dried

using magnesium sulphate and concentrated under reduced pressure. The crude product is stored in the fridge and reacted further.

Example II: 2-Cyclopropyl-3-dimethylaminoacrylonitrile

5



10

At 100°C, 25 g (292.8 mmol) of cyclopropylacetonitrile (obtainable by the process described in US-3,454,575) and 25.5 g (30.2 ml, 146.4 mmol) of bis(dimethylamino)-tert-butyloxymethane are stirred under a riser pipe for 46 hours. Volatile components are evaporated under reduced pressure and the residue is subsequently distilled at 0.1 Torr and 60-65°C.


Yield 16.18 g (81% of theory)

15

In a manner analogous to Examples I and II, it is possible to prepare the corresponding cyclobutyl, cyclopentyl, cyclohexyl and 1-cyclopentene-1-yl derivatives.


Example III: Ethyl 5-amino-1-(2-fluorobenzyl)-1H-pyrazolo-3-carboxylate

20



Under argon, 100 g (0.613 mol) of the sodium salt of ethyl cyanopyruvate (preparation analogously to Borsche and Manteuffel, Liebig's Ann. 1934, 512, 97) in 2.5 l of dioxane are mixed at room temperature and with efficient stirring with 111.75 g (75 ml, 0.98 mol) of trifluoroacetic acid, and the mixture is stirred for 10 min, during which a large fraction of the starting material dissolves. 85.93 g (0.613 mol) of 2-fluorobenzylhydrazine are then added, and the mixture is boiled overnight. After cooling, the precipitated sodium trifluoroacetate crystals are filtered off with suction and washed with dioxane, and the crude solution is reacted further.

10 Example IV: Ethyl 1-(2-fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine-3-carboxylate




The solution obtained in Example III is mixed with 61.25 ml (60.77 g, 0.613 mol) of dimethylaminoacroleine and 56.28 ml (83.88 g, 0.736 mol) of trifluoroacetic acid and boiled under argon for 3 days. The solvent is subsequently evaporated under reduced pressure and the residue is added to 2 l of water and extracted three times with 1 l of ethyl acetate each time. The combined organic phases are dried with magnesium sulphate and concentrated using a rotary evaporator. The residue is chromatographed over 2.5 kg of silica gel, eluting with a toluene/toluene-ethyl acetate = 4:1 gradient.

Yield: 91.6 g (49.9% of theory over two steps).

M.p. 85°C


R_f (SiO₂, TIE1): 0.83

Example V: 1-(2-Fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine-3-carboxamide

- 5 10.18 g (34 mmol) of the ester obtained in Example IV are initially charged in 150 ml of methanol which has been saturated with ammonia at 0 - 10°C. The mixture is stirred at room temperature for two days and subsequently concentrated under reduced pressure.

$R_f(\text{SiO}_2, \text{TIE1}): 0.33$

10

Example VI: 3-Cyano-1-(2-fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine

- 15 36.1 g (133 mmol) of 1-(2-fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine-3-carboxamide from Example V are dissolved in 330 ml of THF and mixed with 27 g (341 mmol) of pyridine. Over a period of 10 min, 47.76 ml (71.66 g, 341 mmol) of trifluoroacetic anhydride are subsequently added, during which the temperature rises to up to 40°C.


The mixture is stirred at room temperature overnight. The mixture is subsequently poured into 1 l of water and extracted three times with 0.5 l of ethyl acetate each time. The organic phase is washed with saturated sodium bicarbonate solution and with 1 N HCl, dried with MgSO_4 and concentrated using a rotary evaporator.

5 Yield: 33.7 g (100% of theory)


M.p.: 81°C

$R_f(\text{SiO}_2, \text{TIE1})$: 0.74

10 Example VII: Methyl 1-(2-fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine-3-carboximidate



30.37 g (562 mmol) of sodium methoxide are dissolved in 1.5 l of methanol, and 36.45 g (144.5 mmol) of 3-cyano-1-(2-fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine (from Example VII) are added. The mixture is stirred at room temperature for 2 hours and the resulting solution is used directly for the next step.


Example VIII: 1-(2-Fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine-3-carboxamide

5 The solution of methyl (2-fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine-3-carboximidate in methanol obtained from Example VII is admixed with 33.76 g (32.19 ml, 562 mmol) of glacial acetic acid and 9.28 g (173 mmol) of ammonium chloride and stirred under reflux overnight. The solvent is evaporated under reduced pressure, the residue is triturated thoroughly with acetone and the precipitated solid is filtered off with suction. The solid is added to 2 l of water, admixed with stirring with
 10 31.8 g of sodium carbonate and extracted three times with a total of 1 l of ethyl acetate, and the organic phase is dried with magnesium sulphate and concentrated under reduced pressure.

Yield: 27.5 g (76.4% of theory over two steps)

M.p.: 86°C

15 $R_f(\text{SiO}_2, \text{TlEtOH})$: 0.08


Example IX: 1-(2-Fluorobenzyl)-3-cyanoindazole

Under argon, 12.0 g (83.9 mmol) of 3-cyanoindazole [for the preparation, cf. Salkowski, Chem. Ber. 17 (1984), 508] were dissolved in 100 ml of abs. THF, and 20.6 g (109 mmol) of 2-fluorobenzyl bromide were added. With ice-cooling, 2.55 g (100 mmol) of sodium hydride (95 per cent) were added a little at a time. The mixture was stirred overnight at room temperature and then concentrated to about one quarter of its volume using a rotary evaporator and admixed with H₂O and ethyl acetate. The aqueous phase was re-extracted with ethyl acetate. The combined organic phases were dried over MgSO₄ and the solvent was distilled off using a rotary evaporator, leaving the product.

Yield: 19.5 g (93%)

R_f value: 0.69 (silica gel; cyclohexane/ethyl acetate 1:1)

Example X: 1-(2-Fluorobenzyl)indazole-3-amidinium chloride




A sodium methoxide solution, which had been prepared from 190 mg (8.26 mmol) of NaOMe and 30 ml of abs. methanol, was added to a solution of 20.0 g (97.7 mmol) of 1-(2-fluorobenzyl)-3-cyanoindazole (from Example IX) in 200 ml of methanol, and the mixture was stirred at 40°C for 22 h. After addition of 0.46 ml of acetic acid and 4.30 g of NH₄Cl, the mixture was stirred at 40°C for another 24 h and subsequently concentrated to dryness using a rotary evaporator. The residue was taken up in acetone and the remaining precipitate was filtered off with suction, giving, after drying under high vacuum, the product in the form of a light-beige powder.

Yield: 20.5 g (84%)

M.p.: >230°C

MS-EI: *m/z*(%) = 268 (31, M⁺ of the free base), 251 (15), 109 (100).

Example XI: 2-Cyanocyclopropylacetamide


5

1.5 g (9.8 mmol) of ethyl 2-cyanocyclopropylacetate (preparation according to Li, J. Med. Chem. 1996, 39, 3070) are dissolved in 14 ml of methanolic ammonia (7N). After four days, the solution is concentrated using a rotary evaporator and the solid is washed with a little methanol.

10

Yield: 456 mg (37.5% of theory).

MS (DCI-NH₃): 142 (100%, M⁺NH₄)

Example XII: 2-Cyclopropylmalononitrile

15

440 mg (3.54 mmol) of 2-cyanocyclopropylacetamide (from Example XI) are admixed twice with toluene which is then evaporated using a rotary evaporator to remove the methanol, and the substance is dissolved in 15 ml of tetrahydrofuran:dichloromethane (2:1). A total of 2.53 g (10.63 mmol) of Burgess reagent is added in three identical portions at intervals of 30 minutes. After a further 30 minutes, the reaction mixture is chromatographed directly over 20 g of silica gel using cyclohexane/ethyl acetate (2:1 to 1:1).

20

Yield: 122 mg (32.4% of theory)

$R_f(\text{SiO}_2, \text{C1E1})$: 0.65

Example XIII: 1-Cyanomethyl-1-fluoromethyl-cyclopropane

5



10

At 140°C, 8.85 g (72.2 mmol) of 1-chloromethyl-1-fluoromethyl-cyclopropane, 4.60 g (93.8 mmol) of sodium cyanide and 433 mg (2.89 mmol) of sodium iodide were heated in 72 ml of triethylene glycol at 140°C for 30 minutes. The mixture was allowed to cool to room temperature and admixed with 150 ml each of dichloromethane and water, and the organic phase was washed with saturated sodium chloride solution, dried over MgSO_4 and concentrated to dryness at room temperature using a rotary evaporator. The liquid product was reacted further without any further purification.

15

Yield: 5.44 g (67%)

Example XIV: 2-[(1-Fluoromethyl)cycloprop-1-yl]-2-formylacetone



20

Under argon, 9.20 g (82.1 mmol) of potassium tert-butoxide were initially charged in 80 ml of anhydrous tetrahydrofuran (THF) and the mixture was mixed with a solution of 4.21 g (37.3 mmol) of 1-cyanomethyl-1-fluoromethyl-cyclopropane (Ex. XIII) and 9.24 g (125 mmol) of ethyl formate in 20 ml of anhydrous THF. The mixture was stirred at room temperature for 3 hours and then concentrated to dryness using a rotary evaporator, and the residue was admixed with 150 ml each of ice water and ethyl acetate. The aqueous phase was separated off and adjusted to pH 4 using

25


dilute hydrochloric acid. It was possible to isolate the product by extracting twice with 100 ml of ethyl acetate each time, drying of the organic phase over MgSO_4 and stripping off the solvent using a rotary evaporator. The product was used without any further purification for the next reaction.

5 Yield: 4.03 g (77%)

09744753-032601

Preparation examples

Example 1: 3-(4-Amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1*H*-pyrazolo[3,4-*b*]pyridine



5

Procedure A

2 g (7.4 mmol) of the amidine from Example VIII are ground until it has a powder-like consistency. The lump-free starting material is mixed with 4 g (29.4 mmol) of 2-cyclopropyl-3-dimethylaminoacrylonitrile from Example II and mixed intimately, using spatular and ultrasonic bath, until a homogeneous milk has formed. Water pump vacuum is applied, causing the mixture to foam. With shaking, the mixture is subsequently immersed into an oil bath at 106°C, and the mixture becomes clear. After 2 hours, the content of the small flask begins to solidify. The mixture is left under reduced pressure at 106°C overnight. The resulting solid is stirred with toluene, filtered off with suction and washed with ether. The residue is taken up in 50 ml of boiling acetonitrile and filtered off with suction. The residue obtained is taken up in 25 ml of boiling dimethylformamide and once more filtered off with suction. Both filtrates are combined and concentrated using a rotary evaporator.

Yield: 0.85 g (31.2% of theory)

M.p.: 210°C

MS (ESI-POS): 361 (100%, M+H)

20

¹H-NMR (300 MHz, d⁶-DMSO): 0.61 (m, 2H, 2-cyclopropyl), 0.9 (m, 2H, 2-cyclopropyl), 1.65 (m, 1H, 1-cyclopropyl), 5.7 (s, 2H, benzyl-CH₂), 6.98 (broad s, 2H, NH₂), 7.1-7.3 (peak cluster, 3H, aromatic benzylic H3,5,6), 7.3-7.4 (peak cluster, 2H, H5, benzylic H4), 8.0 (1H, pyrimidinyl-H6), 8.6 (d, 1H, H6), 8.95 (d, 1H, H4)

5

Procedure B

20.0 g (74.3 mmol) of the amidine from Example VIII and 28.4 g (260.0 mmol) of the crude 2-cyclopropyl-3-oxopropionitrile from Example I are added together in 280 ml of toluene and mixed in an ultrasonic bath, and the mixture is heated at reflux overnight. The solution is concentrated to half its volume using a rotary evaporator, cooled and filtered off with suction, and the filter residue is washed with ethyl acetate. The filtrate is concentrated and chromatographed over silica gel 60 (particle size 0.040-0.063 mm) using cyclohexane/ethyl acetate 1:1 as mobile phase. The residue is taken up in 50 ml of boiling toluene and filtered off with suction. Both solutions are combined, concentrated using a rotary evaporator and recrystallized from toluene.

10

15

Yield: 8.38 g (31.3% of theory).

R_f(SiO₂, C1E2): 0.23

M.p.: 209°C

20

Procedure C (method without solvent)

103.7 mg (0.38 mmol) of the amidine from Example VIII and 168.1 mg (1.54 mmol) of the crude 2-cyclopropyl-3-oxopropionitrile from Example I are added together in 0.5 ml of toluene and mixed in an ultrasonic bath. The toluene is evaporated under reduced pressure and the mixture is heated to 100-105°C without solvent in an open vessel. After one hour, the mixture is cooled, the residue is dissolved in dichloromethane and admixed with 1 g of silica gel and the mixture is concentrated using a rotary evaporator.

25


30

For purification, the substance is chromatographed over silica gel 60 (particle size 0.040-0.063 mm) using cyclohexane/ethyl acetate 1:1 as mobile phase.

Yield: 50.0 mg (36.0% of theory).

$R_f(\text{SiO}_2, \text{C1E2}): 0.23$

The following compounds (Ex. 2 to 5) are prepared analogously to the procedures mentioned above:



Example 2: 3-(4-Amino-5-cyclobutylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine

10 R¹ = cyclobutyl
M.p. 214°C


Example 3: 3-(4-Amino-5-cyclopentylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine

15 R¹ = cyclopentyl
M.p. 208°C

Example 4: 3-(4-Amino-5-cyclohexylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine

20 R¹ = cyclohexyl
M.p. 213°C

Example 5: 3-(4-Amino-5-(1-cyclopenten-1-yl)-pyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine



M.p. 228°C

5

Example 6: 3-(4-Amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine, hydrochloride

10

0.3 g (0.83 mmol) of 3-(4-amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine from Example 1 is dissolved in 50 ml of hot acetonitrile, and 0.9 ml of 1 N HCl (0.9 mmol) is added. The precipitated crystals are subsequently filtered off with suction.

Yield: 0.24 g (72.7% of theory)

M.p. 279°C.

15

Example 7: 3-(4-Amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine, tosylate


20

0.3 g of 3-(4-amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine from Example 1 is dissolved in 50 ml of hot acetonitrile, and 160 mg of p-toluenesulphonic acid are added. After cooling to room temperature, the precipitated crystals are filtered off with suction.

Yield: 303 mg (68.8% of theory),

M.p. 203°C

25




At room temperature, 105 mg (2.62 mmol) of sodium hydride (60% suspension in oil) are suspended in 20 ml of THF. 472 mg (1.31 mmol) of the amine from Example 1 in 10 ml of THF and 432 mg (1.57 mmol) of 2-(benzoyloxymethyl)benzoyl chloride in 5 ml of THF are added. The solution is admixed with water, extracted with ethyl acetate and washed with 1M hydrochloric acid and NaHCO₃ solution, dried with magnesium sulphate and concentrated under reduced pressure. The substance is digested repeatedly with cyclohexane.

Yield: 474 mg (60.4% of theory)

R_f: 0.61 (SiO₂, C1E2)

M.p.: 134-136°C

Example 9: 3-(5-Cyclopropyl-4-(2,2-dimethylpropanoyl)aminopyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine




- 5 60 mg (0.17 mmol) of the amine from Example 1 are taken up in dichloromethane, 30 mg (0.25 mmol) of pivaloyl chloride, 26 mg (0.33 mmol) of pyridine and a catalytic amount of dimethylaminopyridine are added and the solution is stirred at room temperature for four hours. The solution is washed with 1N hydrochloric acid and saturated NaHCO₃ solution, dried with MgSO₄ and concentrated under reduced
- 10 pressure. The crude product is chromatographed over silica gel using cyclohexane/ethyl acetate 1:1 as mobile phase.

Yield: 24.8 mg (33.5% of theory)

Rf: 0.59 (SiO₂, EA)

The following compounds can be obtained in an analogous manner:



5


Ex.	X	Yield (% of theory)	R _F (SiO ₂)
10 (from 1 and CH ₃ COCl)		5.1	0.36(C1E2)
11 (from 1 and CH ₃ COCl)		55.4	0.53(C1E2)
12 (from 1 and CH ₃ SO ₂ Cl)		11.8	0.79(BABA)
13 (from 1 and ethyl chloroformate)		42.9	0.64 (EA)

Ex.	X	Yield (% of theory)	R _f (SiO ₂)
14 (from 1 and the corresponding acyl chloride)		17.0	0.29 (C1E1)
15 (from 1 and 2-chloroethyl chloroformate)		21.3	0.60 (EA)
16 (from 1 and o- azidobenzoyl chloride)		7.1	0.69 (EA)
17 (from 1 and p- azidobenzoyl chloride)		72.9	0.37 (C1E1)
18 (from 1 and m- azidobenzoyl chloride)		26.7	0.32 (C1E1)

BABA: 50 ml of n-butylacetate + 9 ml of n-butanol + 25 ml of glacial acetic acid + 15 ml of phosphate buffer pH 6 are shaken. The aqueous, lower phase that separates off is discarded.

5

Example 19: 3-(5-Cyclopropyl-4-(ethoxymethylamino)pyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine





195.5 mg (0.54 mmol) of the amine from Example 1 are dissolved in 5 ml of ethanol. 0.9 ml (10.85 mmol) of a 37% strength solution of formaldehyde is added, the solution is heated at reflux for 24 hours. The solvents are evaporated under reduced pressure and the crude product is chromatographed over silica gel using cyclohexane/ethyl acetate 2:1 to ethyl acetate.


Yield: 87.1 mg (38.4% of theory)

Rf: 0.51 (SiO₂, EA)

The following compounds can be prepared analogously:


Ex..	X	Yield (% of theory)	R _f (SiO ₂)
20 (from 1 and formaldehyde)		70.9	0.38 (EA)
21 (from 1 and acetaldehyde/ ethanol)		28.0	0.59 (EA)



Example 22: 3-(5-Cyclopropyl-4-(propylaminocarbonylamino)pyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine




- 5 Under argon, 106 mg (0.29 mmol) of the amine from Example 1 are suspended in dichloromethane, and the mixture is admixed with 137.6 mg (1.62 mmol) of propyl isocyanate at 0°C. The mixture is heated at reflux for 38 hours. The substance is chromatographed over silica gel using cyclohexane/ethyl acetate 1:1 to ethyl acetate as mobile phase.
- 10 Yield: 52 mg (39.6% of theory)
Rf: 0.58 (SiO₂, EA)

The following compounds can be prepared analogously:



Ex.	X	Yield (% of theory)	R _f (SiO ₂)
23 (from 1 and m-chlorophenyl isocyanate)		55.8	0.60 (CE)
24 (from 1 and hexyloxymethyl isocyanate)		11.3	0.29 (CE)

Example 25: 3-(4-Amino-5-cyclopropyl-2-pyrimidyl)-1-(2-fluorobenzyl)indazole




- Under argon 1.08 g (6.00 mmol) of a solution of sodium methoxide (30 per cent in methanol) were admixed with 15 ml of abs. methanol and 1.83 g (6.00 mmol) of 1-(2-fluorobenzyl)indazol-3-amidinium chloride from Example X. The mixture was stirred at room temperature for 5 minutes, after which 816 mg (6.00 mmol) of 2-cyclopropyl-3-dimethylaminoacrylonitrile from Example II were added, and the mixture was heated under reflux overnight. After cooling to room temperature, the precipitate was filtered off with suction and stirred in pentane. Once more, the precipitate was filtered off with suction and dried under high vacuum, giving the product in the form of a virtually white solid.

Yield: 700 mg (32%)

M.p.: 218°C

¹H-NMR: (400 MHz, D₆-DMSO), δ = 0.61 (m, 2H, cyclo-Pr-CH₂), 0.91 (m, 2H, cyclo-Pr-CH₂), 1.67 (m, 1H, cyclo-Pr-CH), 5.80 (s, 2H, CH₂), 6.99 (brd. s, 2H, NH₂),
 5 7.04 - 7.13 (m, 2H, Ar-H), 7.20 - 7.27 (m, 2H, Ar-H), 7.31 - 7.38 (m, 1H, Ar-H),
 7.43 (t, 1H, Ar-H), 7.73 (d, 1H, Ar-H), 7.99 (s, 1H, Ar-H), 8.63 (d, 1H, Ar-H).

Example 26: 3-(4,6-Diamino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine







10 155 mg (0.57 mmol) of the amidine from Example VIII and 122 mg (1.15 mmol) of 2-cyclopropylmalonitrile from Ex. XII are added together in dichloromethane and heated in an open vessel to 105°C. After one hour, the reaction mixture solidifies. After three hours, the mixture is cooled, digested with toluene, filtered and dried
 15 under high vacuum.



Yield: 156 mg (72.3% of theory)

R_f(SiO₂, BABA): 0.37


M.p.: 198-200°C

20 Analogously to Example 26, the following compounds are obtained:

Ex.	X	Yield (% of theory)	R _f (SiO ₂)
<p>27</p> <p>(from VIII and</p> 		12.2	0.63 (BABA)
<p>28</p> <p>(from VIII and</p> 		13.2	0.70 (EA)

Ex.	X	Yield (% of theory)	R_f (SiO ₂)
<p>29</p> <p>(from 28 by reaction with POCl₃/PhNMe₂)</p>		100	0.75 (EA)
<p>30</p> <p>(from 28 by reaction with PBr₃/PhNMe₂)</p>		21.9	0.19 (SiO ₂ , C1E1)

Example 31: 3-[4-Amino-5-(1-fluoromethyl)cycloprop-1-yl]-pyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine



- 5 1.00 g (3.70 mmol) of the amidine from Ex. VIII and 1.83 g (13.0 mmol) of 2-[(1-fluoromethyl)cycloprop-1-yl]-2-formylacetonitrile from Ex. XIV were suspended in 30 ml of toluene. The mixture was sonicated with ultrasound for five minutes and subsequently heated under reflux. The mixture was concentrated using a rotary evaporator and then chromatographed over silica gel (C → C1:E1 → EA). The most
- 10 polar fraction had to be subjected to another column chromatography, again using the gradient (C/EA) but with addition of in each case 1% of triethylamine to the mobile phase.


Yield: 486 mg (34% of theory)

M.p.: 220°C

- 15 R_f value: 0.19 (C1:E1)

Patent Claims

1. Substituted pyrazole derivatives of the general formula (I)



in which

at least one of the substituents R^1 , X and Y represents saturated or partially unsaturated C_3 - C_8 -cycloalkyl,

which may optionally be mono- or polysubstituted by amino, azido, formyl, mercaptyl, carboxyl, hydroxyl, morpholino, piperidino, pyrrolidino, sulphonamino, straight-chain, cyclic or branched acyl, acylamino, alkoxy, benzyloxy, alkylamino, dialkylamino, alkylsulphonyl, alkylsulphonamino, alkylthio, alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, halogen, phenyl

and/or is optionally substituted by

straight-chain or branched or cyclic alkyl having up to 6 carbon atoms which for its part may be substituted by amino, mercaptyl, carboxyl, hydroxyl, morpholino, piperidino, pyrrolidino, straight-chain, cyclic or branched acyl, acylamino, alkoxy, alkylamino, dialkylamino, alkylsulphonyl, alkylthio,


phenyl, alkylsulphonamino, alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, halogen,

and where the optionally remaining radicals R^1 , X and/or Y are identical or different and each represents hydrogen, azido, formyl, mercaptyl, carboxyl, hydroxyl, straight-chain or branched acyl, alkoxy, alkylthio or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, halogen, phenyl or straight-chain or branched alkenyl or alkynyl having in each case up to 6 carbon atoms or alkyl having up to 20 carbon atoms, where both alkenyl, alkynyl and/or alkyl may optionally be substituted by hydroxyl, amino, azido, carboxyl, straight-chain or branched acyl, alkoxy, alkoxycarbonyl or acylamino having in each case up to 5 carbon atoms, aryl having 6 to 10 carbon atoms, a 5- to 6-membered aromatic heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, halogen, cyano, dialkylamino having up to 6 carbon atoms, alkylamino having up to 6 carbon atoms and/or cycloalkyl having 3 to 8 carbon atoms or by a radical of the formula $-OR^4$,

in which

R^4 represents straight-chain or branched acyl having up to 5 carbon atoms,

and/or the optionally remaining radicals R^1 , X and/or Y each represents a radical of the formula



in which

5 a, b and b' are identical or different and represent a number 0, 1, 2 or 3,

R⁵ represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

10 and/or the optionally remaining radicals R¹, X and/or Y each represent a 3- to 8-membered ring, which may also be linked to the pyrimidine via a -CO-bridge, and which may be saturated, unsaturated and/or partially unsaturated, which may contain 1 to 4 heteroatoms from the group consisting of N, O, S, SO, SO₂, which may also be attached via N, which may optionally contain a

15 carbonyl group as ring member and which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl, amino, halogen, carboxyl, straight-chain or branched acyl, alkoxy, alkoxy carbonyl or acylamino having

20 in each case up to 5 carbon atoms,

and/or the optionally remaining radicals R¹, X and/or Y each represent straight-chain or branched acyl having up to 6 carbon atoms which is optionally substituted by halogen,

25 or

represent straight-chain or branched acyloxy having up to 6 carbon atoms,

or

represent arylthio having 6 to 10 carbon atoms or heteroarylthio,

and/or the optionally remaining radicals R^1 , X and/or Y represent radicals of the formulae $-SO_3H$ or $S(O)_2R^6$,

in which

c represents a number 1 or 2,

R^6 represents straight-chain or branched alkyl having 1 to 10 carbon atoms, cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or a 5- to 6-membered heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, where the ring systems may optionally be substituted by halogen or by straight-chain or branched alkyl or alkoxy having in each case up to 4 carbon atoms,

and/or the optionally remaining radicals R^1 , X and/or Y each represent a radical of the formula $PO(OR^7)(OR^8)$,

in which

R^7 and R^8 are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 8 carbon atoms or cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or benzyl,

and/or the optionally remaining radicals R^1 , X and/or Y each represent oxycycloalkyl having 3 to 8 carbon atoms or represent radicals of the formulae

$-\text{NH}-\text{C}(=\text{NH})\text{NH}_2$, $\text{CON}=\text{C}(\text{NH}_2)_2$ or $-\text{C}=\text{NH}(\text{NH}_2)$, $(\text{CO})_d\text{NR}^9\text{R}^{10}$ or $-\text{NHCONR}^9\text{R}^{10}$,


in which

d represents a number 0 or 1,

R^9 and R^{10} are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 14 carbon atoms or cycloalkyl having 3 to 14 carbon atoms, aryl having 6 to 10 carbon atoms or a 3- to 10-membered ring having up to 5 heteroatoms from the group consisting of N, O, S, which may also be attached via N, where the abovementioned radicals may optionally be substituted by aryl having 6 to 10 carbon atoms, heterocyclyl, cycloalkyl having 3 to 7 carbon atoms, hydroxyl, amino or straight-chain or branched alkoxy, acyl or alkoxycarbonyl having in each case up to 6 carbon atoms,

and in the case that $d = 0$,

R^9 and R^{10} may also represent straight-chain, branched or cyclic acyl having up to 14 carbon atoms, straight-chain or branched hydroxyalkyl having up to 6 carbon atoms, straight-chain or branched alkoxyalkyl having a total of up to 12 carbon atoms, straight-chain or branched alkoxycarbonyl or acyloxyalkyl having in each case up to 6 carbon atoms or a radical of the formula $-\text{SO}_2\text{R}^{11}$ or radicals of the formulae








in which




R^{11} represents straight-chain or branched alkyl having up to 4 carbon atoms,

5

and/or

R^9 and R^{10} represent radicals of the formulae





in which

R^{12} , R^{13} and R^{15} to R^{28} are identical or different and each represents hydrogen
or straight-chain or branched alkyl having up to 4 carbon atoms,

e represents a number 0, 1 or 2,

and

R^{14} represents straight-chain or branched alkyl having up to 6 carbon
atoms or cycloalkyl having 3 to 8 carbon atoms,

R^9 and R^{10} are identical or different and each represents hydrogen; alkyl
having up to 14 carbon atoms which is optionally substituted by
hydroxyl or alkoxy having up to 8 carbon atoms; aryl having 6 to 10
carbon atoms which is optionally substituted by halogen,

R^2 and R^3 form, together with the double bond, a fused phenyl ring or a fused 6-membered saturated or aromatic heterocycle having up to 3 heteroatoms from the group consisting of N, S and O,

5 which is optionally substituted up to 3 times by identical or different substituents from the group consisting of formyl, carboxyl, hydroxyl, mercaptyl, straight-chain or branched acyl, alkylthio or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, cyano, halogen or straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms
10 which for its part may be substituted by hydroxyl, amino, carboxyl, straight-chain or branched acyl, alkoxy or alkoxycarbonyl having in each case up to 5 carbon atoms,

and/or the fused phenyl ring or the fused 6-membered saturated or aromatic heterocycle is optionally substituted by a group of the formula $-NR^{29}R^{30}$,
15

in which

R^{29} and R^{30} are identical or different and each represents hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,
20

or

R^{29} represents hydrogen and
25

R^{30} represents acyl having up to four carbon atoms

and/or the fused phenyl ring or fused 6-membered saturated or aromatic heterocycle are optionally substituted by phenyl which for its part may be substituted up to 2 times by identical or different substituents from the group
30

consisting of halogen, straight-chain or branched alkyl and alkoxy having in each case up to 6 carbon atoms

5 and/or the fused phenyl ring or the fused 6-membered saturated or aromatic heterocycle are optionally substituted by a group of the formula - $\text{N}=\text{CH}-\text{NR}^{31}\text{R}^{32}$,

in which

10 R^{31} and R^{32} are identical or different and each represents hydrogen, phenyl or straight-chain or branched alkyl having up to 6 carbon atoms,

A represents a 5- or 6-membered aromatic or saturated heterocycle
15 having up to 3 heteroatoms from the group consisting of S, N and O or represents phenyl,

which are optionally substituted up to 3 times by identical or different substituents from the group consisting of amino, mercaptyl, hydroxyl, formyl, carboxyl, straight-chain or branched acyl, alkylthio, alkyloxyacyl, alkoxy or
20 alkoxy carbonyl having in each case up to 6 carbon atoms, nitro, cyano, trifluoromethyl, azido, halogen, phenyl and straight-chain or branched alkyl having up to 6 carbon atoms which for its part may be substituted by hydroxyl, carboxyl, straight-chain or branched acyl, alkoxy or alkoxy carbonyl having in each case up to 5 carbon atoms,

25 and/or is substituted by a group of the formula $-(\text{CO})_f-\text{NR}^{33}\text{R}^{34}$,

in which

30 f represents a number 0 or 1,

R^{33} and R^{34} are identical or different and represent hydrogen, phenyl, benzyl or straight-chain or branched alkyl or acyl having in each case up to 5 carbon atoms,

5 and their isomeric forms and salts.

2. Compounds according to Claim 1,

in which

10

at least one of the substituents R^1 , X and Y represents cyclopropyl, cyclobutyl, cyclopentenyl, cyclohexyl or cycloheptyl which may optionally be mono- or polysubstituted by amino, azido, formyl, mercaptyl, carboxyl, hydroxyl, morpholino, piperidino, pyrrolidino, sulphonamino, straight-chain, 15 cyclic or branched acyl, acylamino, alkoxy, benzyloxy, alkylamino, dialkylamino, alkylsulphonyl, alkylsulphonamino, alkylthio, alkoxycarbonyl having in each case up to 4 carbon atoms, nitro, cyano, halogen, phenyl and/or is optionally substituted by

20

straight-chain or branched or cyclic alkyl having up to 4 carbon atoms which for its part may be substituted by amino, mercaptyl, carboxyl, hydroxyl, morpholino, piperidino, pyrrolidino, straight-chain, cyclic or branched acyl, acylamino, alkoxy, alkylamino, dialkylamino, alkylsulphonyl, alkylthio, phenyl, alkylsulphonamino, alkoxycarbonyl having in each case up to 4 carbon atoms, nitro, cyano, halogen,

25

and where the optionally remaining radicals R^1 , X and/or Y are identical or different and each represents hydrogen, azido, formyl, mercaptyl, carboxyl, hydroxyl, straight-chain or branched acyl, alkoxy, alkylthio or alkoxycarbonyl having in each case up to 4 carbon atoms, nitro, cyano, halogen, phenyl or straight-chain or branched alkenyl or alkynyl having in 30 each case up to 4 carbon atoms or alkyl having up to 18 carbon atoms, where both alkenyl, alkynyl and/or alkyl may optionally be substituted by hydroxyl,


30

amino, azido, carboxyl, straight-chain or branched acyl, alkoxy, alkoxy carbonyl or acylamino having in each case up to 4 carbon atoms, phenyl, naphthyl or pyridyl, halogen, cyano, dialkylamino having up to 6 carbon atoms, alkylamino having up to 4 carbon atoms and/or cyclopropyl, cyclopentyl, cyclohexyl or by a radical of the formula $-OR^4$,

in which

R^4 represents straight-chain or branched acyl having up to 4 carbon atoms,

and/or the optionally remaining radicals R^1 , X and/or Y each represent a radical of the formula



in which

a, b and b' are identical or different and represent a number 0, 1 or 2,

and/or the optionally remaining radicals R^1 , X and/or Y each represent a 3- to 8-membered ring, which may also be linked to the pyrimidine via a -CO-bridge, and which may be saturated, unsaturated and/or partially unsaturated, which may contain 1 to 3 heteroatoms from the group consisting of N, O, S, SO, SO₂, which may also be attached via N, which may optionally contain a carbonyl group as ring member and which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of straight-chain or branched alkyl having up to 4 carbon atoms which is optionally substituted by hydroxyl, amino, halogen, carboxyl,

straight-chain or branched acyl, alkoxy, alkoxycarbonyl or acylamino having in each case up to 4 carbon atoms,

5 and/or the optionally remaining radicals R^1 , X and/or Y each represent straight-chain or branched acyl having up to 4 carbon atoms which is optionally substituted by halogen,

or

10 represent straight-chain or branched acyloxy having up to 4 carbon atoms,

or

15 represents phenylthio,

and/or

represent radicals of the formulae $-SO_3H$ or $S(O)_cR^6$,

20 in which

c represents a number 1 or 2,

25 R^6 represents straight-chain or branched alkyl having 1 to 8 carbon atoms, phenyl or a 5- to 6-membered heterocycle having up to 2 heteroatoms from the group consisting of S, N and/or O, where the ring systems may optionally be substituted by halogen or by straight-chain or branched alkyl or alkoxy having in each case up to 3 carbon atoms,

30

and/or the optionally remaining radicals R^1 , X and/or Y each represent a radical of the formula $PO(OR^7)(OR^8)$,

in which

R^7 and R^8 are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 4 carbon atoms or cyclopropyl, cyclopentyl, phenyl or benzyl,

and/or the optionally remaining radicals R^1 , X and/or Y each represent oxycycloalkyl having 3 to 6 carbon atoms or represent radicals of the formulae

$-NH-C(=NH)NH_2$, $-CON=C(NH_2)_2$ or $-C=NH(NH_2)$, $(CO)_dNR^9R^{10}$ or $-NHCONR^9R^{10}$,

in which


d represents a number 0 or 1,

R^9 and R^{10} are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 4 carbon atoms or cyclopropyl, cyclopentyl, cyclohexyl, phenyl or a 3- to 6-membered ring having up to 3 heteroatoms from the group consisting of N, O, S, which may also be attached via N,

where the abovementioned radicals may optionally be substituted by phenyl, cyclopropyl, cyclopentyl, hydroxyl, amino or straight-chain or branched alkoxy, acyl or alkoxy-carbonyl having in each case up to 4 carbon atoms,

and in the case that $d = 0$,

R^9 and R^{10} may also represent straight-chain, branched or cyclic acyl having up to 6 carbon atoms, straight-chain or branched hydroxyalkyl having up to 4 carbon atoms, straight-chain or branched alkoxyalkyl having a total of up to 10 carbon atoms, straight-chain or branched alkoxycarbonyl or acyloxyalkyl having in each case up to 4 carbon atoms or a radical of the formula $-SO_2R^{11}$ or a radical of the formulae










in which

R^{11} represents straight-chain or branched alkyl having up to 3 carbon atoms,

and/or

R^9 and R^{10} represent radicals of the formulae





in which

R^{12} , R^{13} and R^{15} to R^{28} are identical or different and each represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

e represents a number 0, 1 or 2,

and

R^{14} represents straight-chain or branched alkyl having up to 6 carbon atoms or cycloalkyl having 3 to 8 carbon atoms,

R^9 and R^{10} are identical or different and each represents hydrogen; alkyl having up to 4 carbon atoms which is optionally substituted by hydroxyl or alkoxy having up to 7 carbon atoms, phenyl which is optionally substituted by halogen,

R^2 and R^3 , together with the double bond, form a fused phenyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl ring,

which are optionally substituted up to 2 times by identical or different substituents from the group consisting of formyl, carboxyl, hydroxyl, mercaptyl, straight-chain or branched acyl, alkylthio or alkoxycarbonyl having in each case up to 5 carbon atoms, nitro, cyano, azido, fluorine, chlorine, bromine or straight-chain or branched alkyl or alkoxy having in each case up to 5 carbon atoms which for its part may be substituted by hydroxyl, amino, carboxyl, straight-chain or branched acyl, alkoxy or alkoxycarbonyl having in each case up to 4 carbon atoms,

and/or

the abovementioned heterocyclic rings or phenyl are optionally substituted by a group of the formula $-NR^{29}R^{30}$,

in which

R²⁹ and R³⁰ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms

5 or

R²⁹ represents hydrogen

and

10

R³⁰ represents formyl

and/or the abovementioned fused phenyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl rings are optionally substituted by phenyl which for its part may be substituted by fluorine, chlorine, bromine or by straight-chain or branched alkyl or alkoxy having in each case up to 4 carbon atoms,

15

A represents thienyl, tetrahydropyranyl, tetrahydrofuranyl, phenyl, morpholinyl, pyrimidyl, pyrazinyl, pyridazinyl or pyridyl which are optionally substituted up to 2 times by identical or different substituents from the group consisting of hydroxyl, formyl, carboxyl, straight-chain or branched acyl, alkylthio, alkylloxyacyl, alkoxy or alkoxycarbonyl having in each case up to 4 carbon atoms, fluorine, chlorine or bromine,

20

25 and their isomeric forms and salts.

3. Compounds according to Claim 1,

in which

30

at least one of the substituents R^1 , X and Y represents cyclopropyl which is optionally substituted by hydroxyl or fluoromethyl, or represents cyclobutyl, cyclopentenyl, cyclopentyl or cyclohexyl,

5 and where the optionally remaining radicals R^1 , X and/or Y are identical or different and each represents hydrogen, hydroxyl, halogen or azido,

and/or

10 represent a 3- to 6-membered ring which may be saturated, unsaturated and/or partially unsaturated and may contain 1 to 3 heteroatoms from the group consisting of N, O, S, SO, SO_2 , which may optionally contain a carbonyl group as ring member, which may also be attached via N and which is
15 the group consisting of straight-chain or branched alkyl having up to 4 carbon atoms which is optionally substituted by hydroxyl, amino, halogen, carboxyl, straight-chain or branched acyl, alkoxy, alkoxycarbonyl or acylamino having in each case up to 4 carbon atoms,

20 and/or

represent straight-chain or branched acyl having up to 4 carbon atoms which is optionally substituted by halogen,

25 or

represent straight-chain or branched acyloxy having up to 4 carbon atoms,

and/or

30 represent radicals of the formulae $-SO_3H$ or $S(O)_2R^6$,

in which

c represents a number 1 or 2,

R⁶ represents straight-chain or branched alkyl having 1 to 8 carbon atoms, phenyl or a 5- to 6-membered heterocycle having up to 2 heteroatoms from the group consisting of S, N and O, where the ring systems may optionally be substituted by halogen or by straight-chain or branched alkyl or alkoxy having in each case up to 3 carbon atoms,

and/or

represent a radical of the formula PO(OR⁷)(OR⁸),

in which

R⁷ and R⁸ are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 4 carbon atoms or cyclopropyl, cyclopentyl, phenyl or benzyl,

and/or

represent oxycycloalkyl having 3 to 6 carbon atoms or represent radicals of the formulae -CON=C(NH₂)₂ or -C=NH(NH₂) or (CO)_dNR⁹R¹⁰ or NHCONR¹²R¹³,


in which

d represents a number 0 or 1,

R^9 and R^{10} are identical or different and each represents hydrogen, straight-chain or branched alkyl having up to 4 carbon atoms or cyclopropyl, cyclopentyl, cyclohexyl or phenyl,

and in the case where $d = 0$

R^9 and R^{10} also represent straight-chain, branched or cyclic acyl having up to 5 carbon atoms, straight-chain or branched hydroxyalkyl having up to 3 carbon atoms, straight-chain or branched alkoxyalkyl having a total of up to 8 carbon atoms, straight-chain or branched alkoxy carbonyl or acyloxyalkyl having in each case up to 3 carbon atoms or a radical of the formula $-SO_2R^{11}$ or a radical of the formulae











in which

R^{11} represents straight-chain or branched alkyl having up to 4 carbon atoms

and/or

R^9 and R^{10} represent radicals of the formulae





in which

R^{12} , R^{13} and R^{15} to R^{28} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

e represents a number 0, 1 or 2

and

R^{14} represents straight-chain or branched alkyl having up to 6 carbon atoms or cycloalkyl having 3 to 8 carbon atoms,

R^{29} and R^{10} are identical or different and represent hydrogen; alkyl having up to 4 carbon atoms which is optionally substituted by hydroxyl or alkoxy having up to 7 carbon atoms, phenyl which is optionally substituted by halogen,

R^2 and R^3 form, together with the double bond, a phenyl, pyridyl or pyrimidinyl ring,


A represents phenyl or pyrimidyl, each of which is optionally substituted by fluorine, chlorine or bromine,

and their isomeric forms and salts.

4. Compounds according to any of the preceding claims,

in which

- 5 R^1 , X and Y are attached to the pyrimidine ring as follows



and

10

R^1 represents an optionally substituted cyclopropyl, cyclobutyl, cyclopentenyl, cyclopentyl, cyclohexyl, 1-hydroxycyclopropyl or 1-(fluoromethyl)cyclopropyl radical,

15

X represents NH_2

and

Y represents hydrogen or NH_2 .

20


5. Compounds according to claim 4, in which R^1 represents an optionally substituted cyclopropyl radical.

25

6. Process for preparing the compounds of the general formula (I) according to Claim 1, characterized in that,

depending on the various meanings of the heterocycles listed above under R^2 and R^3 ,

[A] compounds of the general formula (II)




in which

10 R^1 , X and Y are each as defined above


and

15 D represents radicals of the formulae



20 R^{38} represents C_1 - C_4 -alkyl


are converted, by reaction with compounds of the general formula (III)



in which

A is as defined above

in inert solvents, if appropriate in the presence of a base, into the compounds of the general formula (IV) or (IVa)



in which


A, X, Y and R^1 are each as defined above,

and, in the case of the compounds of the general formula (IVa), are subsequently cyclized with carboxylic acids, nitriles, formamides or guanidium salts,

and, in the case of the compounds of the general formula (IV), are cyclized with 1,3-dicarbonyl derivatives, their salts, tautomers, enol ethers or enamines, in the presence of acids and, if appropriate, under microwave irradiation,

or


[B] in the case that R^2 and R^3 together form a pyrazine ring, compounds of the general formula (IV) are initially converted by nitrosation into the compounds of the general formula (V)



in which

A, X, Y and R^1 are each as defined above,

in a second step, the compounds of the general formula (VI)



in which


A, X, Y and R^1 are each as defined above

are prepared by a reduction,

and these are subsequently cyclized with 1,2-dicarbonyl compounds, preferably aqueous glyoxal solution,

or

[C] compounds of the general formula (VII)



in which

A¹, R² and R³ are each as defined above

and

L represents a radical of the formula -SnR³⁶R³⁷R³⁸, ZnR³⁹, iodine, bromine or triflate,


in which

R³⁶, R³⁷ and R³⁸ are identical or different and each represents straight-chain or branched alkyl having up to 4 carbon atoms

and

R³⁹ represents halogen

are reacted with compounds of the general formula (VIII)



(VIII),

in which

X, Y and R¹ are each as defined above

and

in the case that L = SnR³⁶R³⁷R³⁸ or ZnR³⁹,

T represents triflate or represents halogen, preferably bromine

and,

in the case that L = iodine, bromine or triflate,

T represents a radical of the formula SnR³⁶R³⁷R³⁸, ZnR³⁹ or BR⁴⁰R⁴¹,

in which


R³⁶, R³⁷, R³⁸ and R³⁹ have the meanings of R³⁶, R³⁷, R³⁸ and R³⁹ given above and are identical to or different from them,

R⁴⁰ and R⁴¹ are identical or different and each represent hydroxyl, aryloxy having 6 to 10 carbon atoms or straight-chain or branched alkyl or alkoxy having in each case up to 5 carbon atoms, or together form a 5- or 6-membered carbocyclic ring

in a palladium-catalysed reaction in inert solvents, if appropriate in the presence of a base,

or

5 [D] amidines of the general formula (IX)




in which

10

A, R² and R³ are each as defined above

are reacted, for example, with compounds of the general formula (X), (Xa),
(Xb) or (Xc)


15




(X)



(Xa)



(Xb)



(Xc)

in which

R^{1'} represents the optionally substituted cycloalkyl radical listed above under R¹;


Alk represents straight-chain or branched alkyl having up to 8 carbon atoms, preferably up to four carbon atoms;

and

Z represents an NH₂ group, a monoalkylamino group having up to 7 carbon atoms, a dialkylamino group having up to 7 carbon atoms, a piperidinyl or morpholinyl radical which is attached via the nitrogen, hydroxyl, alkoxy having up to 7 carbon atoms, acyloxy having up to 7 carbon atoms or aryloxy having 6 to 10 carbon atoms,

and, if appropriate, the substituents listed under X, Y, R¹, R², R³ and/or A are modified or introduced by customary methods, preferably by acylation and derivatization of free amino groups, chlorination, catalytic hydrogenation, reduction, oxidation, removal of protective groups and/or nucleophilic substitution.

7. Amidines of the general formula (IX)



(IX),

in which

R², R³ and A are each as defined in one of the preceding Claims 1 to 3,

and their isomeric forms and salts.

- 5 8. Medicaments, comprising at least one compound of the general formula (I) according to Claim 1.
- 10 9. Process for preparing medicaments, characterized in that at least one compound of the formula (I) according to Claim 1, if appropriate with customary auxiliaries and additives, is converted into a suitable administration form.
- 15 10. Medicaments, comprising at least one compound of the general formula (I) according to Claim 1 in combination with organic nitrates or NO donors.
- 20 11. Medicaments, comprising at least one compound of the general formula (I) according to Claim 1 in combination with compounds which inhibit the degradation of cyclic guanosine monophosphate (cGMP).
- 25 12. Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments.
- 30 13. Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments for the treatment of cardiovascular diseases.
14. Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments for the treatment of hypertension.
15. Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments for the treatment of thromboembolic disorders and ischaemia.
16. Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments for the treatment of sexual dysfunction.


17. Use of compounds of the general formula (I) according to Claim 1 for preparing medicaments having anti-inflammatory properties.
- 5 18. Use according to any of Claims 12 to 17 where the compounds of the general formula (I) according to Claim 1 are used in combination with organic nitrates or NO donors or in combination with compounds which inhibit the degradation of cyclic guanosine monophosphate (cGMP).

09744703.000001

Substituted pyrazole derivatives

Abstract

The present invention relates to novel substituted pyrazole derivatives of the general formula (I)



in which the substituents R¹, R², R³, A, X and Y are each as defined, and to processes for their preparation and to their use as medicaments, in particular as medicaments for the treatment of cardiovascular disorders.

FIG. 1

Stimulation of soluble guanylate cyclase by 3-(4-amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]-pyridine (Ex. 1) in the presence of different NO concentrations





FIG. 2

Effect of 3-(4-amino-5-cyclopropylpyrimidin-2-yl)-1-(2-fluorobenzyl)-1H-pyrazolo[3,4-b]pyridine (Ex. 1) on the average blood pressure of awake, spontaneously hypertensive rats

5



COMBINED DECLARATION AND POWER OF ATTORNEY

ATTORNEY DOCKET NO

Le A 33 187

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought

on the invention entitled

SUBSTITUTED PYRAZOLE DERIVATES

the specification of which is attached hereto,

or was filed on **July 16, 1999**

as a PCT Application Serial No. **PCT/EP99/05073**

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s), the priority(ies) of which is/are to be claimed:

198 34 047.8
(Number)

Germany
(Country)

July 29, 1998
(Month/Day/Year Filed)

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose the material information as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)

(Filing Date)

(Status)

(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

(Status)

(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Le A 33 187-US

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and this application and to transact all business in the Patent and Trademark Office connected with this application.

Jeffrey M. Greenman, Reg.No. 26552
Barbara A. Shimei, Reg. No. 29862
William F. Gray, Reg. No. 31018
Alice A. Brewer, Reg. No. 32888
Jerrie L. Chiu, Reg. No. 41670

all of Bayer Corporation, 400 Morgan Lane, West Haven, Connecticut 06516

Send Correspondence To: Direct Telephone Calls To:
Mr. Jeffrey M. Greenman
Bayer Corporation (203)812-3964 (Jerrie L. Chiu)
400 Morgan Lane
West Haven, Connecticut 06516

1-00 FULL NAME OF SOLE OR FIRST INVENTOR <u>Alexander Straub</u>		INVENTOR'S SIGNATURE <u>Alexander Straub</u>	DATE 5.1.2001
RESIDENCE D 42113 Wuppertal, Germany		DEX	CITIZENSHIP German
POST OFFICE ADDRESS c/o Bayer Aktiengesellschaft, D 51368 Leverkusen, Germany			
2-00 FULL NAME OF SECOND INVENTOR <u>Achim Feuerer</u>		INVENTOR'S SIGNATURE <u>Achim Feuerer</u>	DATE 11.12.2000
RESIDENCE D 51519 Odenthal, Germany		DEX	CITIZENSHIP German
POST OFFICE ADDRESS c/o Bayer Aktiengesellschaft, D 51368 Leverkusen, Germany			
3-00 FULL NAME OF THIRD INVENTOR <u>Cristina Alonso-Alija</u>		INVENTOR'S SIGNATURE <u>Cristina Alonso-Alija</u>	DATE 6.12.00
RESIDENCE D 42781 Haan, Germany		DEX	CITIZENSHIP Spanish
POST OFFICE ADDRESS c/o Bayer Aktiengesellschaft, D 51368 Leverkusen, Germany			
4-00 FULL NAME OF FOURTH INVENTOR <u>Elke Stahl</u>		INVENTOR'S SIGNATURE <u>Elke Stahl</u>	DATE 18.12.00
RESIDENCE D 51467 Bergisch Gladbach, Germany		DEX	CITIZENSHIP German
POST OFFICE ADDRESS c/o Bayer Aktiengesellschaft, D 51368 Leverkusen, Germany			
5-00 FULL NAME OF FIFTH INVENTOR <u>Johannes-Peter Stasch</u>		INVENTOR'S SIGNATURE <u>Johannes-Peter Stasch</u>	DATE 12.12.2000
RESIDENCE D 42651 Solingen, Germany		DEX	CITIZENSHIP German
POST OFFICE ADDRESS c/o Bayer Aktiengesellschaft, D 51368 Leverkusen, Germany			
6-00 FULL NAME OF SIXTH INVENTOR <u>Elisabeth Perzborn</u>		INVENTOR'S SIGNATURE <u>Elisabeth Perzborn</u>	DATE 11.12.2000
RESIDENCE D 42327 Wuppertal, Germany		DEX	CITIZENSHIP German
POST OFFICE ADDRESS c/o Bayer Aktiengesellschaft, D 51368 Leverkusen, Germany			
7-00 FULL NAME OF SEVENTH INVENTOR <u>Joachim Hütter</u>		INVENTOR'S SIGNATURE <u>Joachim Hütter</u>	DATE 18.12.2000
RESIDENCE D 42349 Wuppertal, Germany		DEX	CITIZENSHIP German
POST OFFICE ADDRESS c/o Bayer Aktiengesellschaft, D 51368 Leverkusen, Germany			

FULL NAME OF EIGHTH INVENTOR Klaus Dembowski		INVENTOR'S SIGNATURE <i>Klaus Dembowski</i>		DATE 11.12.2008
RESIDENCE D 69198 Schriesheim, Germany				
POST OFFICE ADDRESS c/o Bayer Aktiengesellschaft, D 51368 Leverkusen, Germany				
FULL NAME OF NINTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				
FULL NAME OF TENTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				
FULL NAME OF ELEVENTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				
FULL NAME OF TWELFTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				
FULL NAME OF THIRTEENTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				
FULL NAME OF FOURTEENTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				
FULL NAME OF FIFTEENTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				
FULL NAME OF SIXTEENTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				
FULL NAME OF SEVENTEENTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE			CITIZENSHIP	
POST OFFICE ADDRESS				